



# Avian Influenza

Last update: 2025-07-04

## Key facts

- Avian influenza, or bird flu, is a viral infection caused by influenza type A viruses. Notable strains include H5N1 and H7N9, which can cause severe illness in humans.
- There are many different influenza (flu) viruses, and they often circulate at the same time. Also, some are seasonal, circulating among humans, while others may be zoonotic, transmitted from birds and pigs. Avian influenza, or "bird flu", is the influenza virus that affects birds. Some avian influenza viruses have developed the ability to infect other animals too. Very rarely, an avian flu virus will infect a person who has very close contact with birds.
- Usually, humans cannot spread avian flu from one person to another. However, there is a small chance the virus can adapt in such a way that it becomes easy for people to spread the virus from person to person. This is dangerous because people would not have any immunity to this virus as they have never been exposed to it before. It could spread quickly and the whole population would be vulnerable to getting sick. This makes avian influenza of eminent pandemic potential.
- Avian Influenza (especially the strains of Highly Pathogenic Avian Influenza [HPAI]) often has devastating economic losses and consequences for the poultry industry, farmer's livelihoods, international trade, and the health of wild birds.

## Transmission

- Direct contact or handling an infected bird (or other infected animals, especially those that eat wild birds)
  - For example, if a person touches an infected bird and then touches their own nose, mouth or eyes, the virus can spread to the person
- Indirect contact with the environment that has been contaminated with bodily fluids, feathers or eggs from an infected bird(s) (or other infected animal).
- Though it is not common, there is the chance the virus could mutate once a human has been affected. If this happens, human-to-human transmission (same as seasonal influenza transmission) is possible.
- Contact with infected wild birds (or other infected animal)
- Ingesting infected under-cooked poultry products.

## Most Vulnerable to contracting the disease:

- People who are in close and repeated contact with infected birds (or other infected animals) or heavily contaminated environments are at risk for acquiring avian influenza. These may include:
  - Poultry farmers and workers: High exposure risk due to close contact with birds.
  - Veterinarians and animal health workers: Risk from handling and treating birds.
  - Residents near poultry farms/markets: Increased risk due to proximity to infected birds.
  - Healthcare workers: Risk from treating infected patients, especially without PPE.
- Due to outbreaks of various subtypes of the avian influenza virus which increases the risk of mutation and spread to humans, the disease continues to be a global public health concern

## Signs and Symptoms

### In birds:

- sudden death with little to no clinical signs
- Discharge from the nose and eyes,
- coughing,
- Difficulty in breathing or shortness of breath
- swelling of the sinuses and/or head,
- apathy,
- reduced vocalisation,
- marked reduction in feed and water intake,
- cyanosis of the unfeathered skin, wattles and comb
- incoordination and nervous signs and diarrhoea

### In humans:

- Mild to moderate
  - Fever
  - Cough
  - Sore throat
  - Muscle aches
  - Inflammation and redness of the eyes
- Severe
  - Pneumonia
  - Acute respiratory distress syndrome (ARDS) - causing extreme difficulty in breathing
  - Failure of several organs
  - Death

# What can you do to prevent and control an epidemic?

## Monitoring the community and identifying birds and people

- Identify and isolate sick birds and people before they spread the disease to others
- Investigation must be done in both animal and human populations, and health workers in both sectors should work together and share information.
- People should stay away from areas known to be affected by animal influenza viruses, including farms, contaminated areas, sick or dead animals, wild birds, and settings where live animals may be sold or slaughtered
- The public should strictly avoid contact with sick or dead animals of unknown causes, including wild birds. Identify and report dead animals or request their removal by contacting local wildlife or veterinary authorities
- Advise people to stay away from areas known to be affected by avian influenza (including farms, contaminated areas, sick or dead birds, wild birds, and settings where live birds may be sold or slaughtered)

## Treatment and management

- Referral of serious cases
  - In birds; sudden death of no apparent causes to animal health professionals
  - In humans; (example: high fever, difficulty breathing, rapid respirations) to health facilities

## Safe shelters and spaces

- Reduce overcrowding and improve ventilation in living shelters, workplaces, and schools if possible

## Hand and respiratory hygiene

- Promote good hand hygiene (handwashing with soap)
- Promote respiratory hygiene and coughing etiquette (cover your cough or sneeze using your sleeve or a tissue, wash hands after coughing or sneezing, do not spit onto the ground or in public)
- Use personal protection (for example, face mask)

## Social mobilization and behaviour change

- Find out the specific advice being given by health and other relevant authorities in animal health

- Promote recommended health practices (such as social distancing)
- Model following this advice and inform community members of current health practice advice
- Offer support and encouragement to follow the advice
  - Try to gain understanding about if and why health practice advice is not being followed
  - With the guidance of your supervisor and health authorities, work with communities to overcome barriers to following health advice and recommended practices

## Mapping and community assessment

- Make a map of the community.
- Mark the following information on the map:
  - How many people and birds have fallen sick with avian influenza? Where?
  - How many people and birds have died? Where? When?
  - Who and where are the vulnerable people?
  - Where are the local health facilities and services? (include traditional healers)
  - Where are the local poultry farms or bird markets? (include markets that sell live and dead birds)
- Record the following information on the back of the map:
  - When did people start to fall sick with avian influenza?
  - How many people live in the affected community?
    - How many people work directly with birds on poultry farms, in markets, etc?
  - How common is it for people to live together in crowded spaces? Is there ventilation and fresh air in homes, schools and workplaces?
  - Are children badly affected by avian influenza? Are there other groups (specific ages, occupations, geographic areas, etc.) that are badly affected?
  - Do any animal health agencies, veterinarians or agriculture ministry agencies work in the area?
  - What are the community's habits, practices and beliefs about caring for and feeding sick people? Consider any differences in roles and responsibilities between women and men.
    - When babies and infants are sick, do women continue to breastfeed them?
  - Is a social mobilization or health promotion programme in place?
  - Which sources do people use/trust the most for information?
    - Are there rumours or misinformation about avian influenza? What are the rumours

## Volunteer actions

- [01. Community-based surveillance](#)
- [02. Community mapping](#)
- [03. Communicating with the community](#)
- [05. Volunteer protection and safety](#)

- [06. Personal protection equipment \(PPE\) for highly infectious diseases](#)
- [20. Isolating sick people](#)
- [26. Respiratory hygiene and coughing etiquette](#)
- [27. Shelter and ventilation](#)
- [29. Hygiene promotion](#)
- [34. Handwashing with soap](#)
- [35. Handwashing in a highly infectious epidemic](#)
- [43. Social mobilization and behaviour change](#)
- [44. Dealing with rumors](#)

## Other resources

- Centers for Disease Control and Prevention (CDC); [Prevention and Antiviral Treatment of Avian Influenza A Viruses in People](#) (2024)
- Centers for Disease Control and Prevention (CDC); [Avian Influenza in Birds: Causes and How It Spreads](#) (2024)
- World Health Organization (WHO); [Influenza: Avian](#) (2024)
- World Organisation for Animal Health (WOAH); [Avian Influenza](#) (n.d.)
- World Organisation for Animal Health (WOAH); [Avian Influenza](#) (2009)

# 01. Community-based surveillance

## Overview

- Community-based surveillance is the systematic detection and reporting of significant public health events (such as sudden illness or death in people or animals) within a community by community members and volunteers. It is a simple, adaptable, low-cost public health initiative designed to complement early warning systems for potential epidemic diseases.
- Volunteers use something called a “community-case definition” to detect and report signs and symptoms of potential diseases, health risks and events, and support in community actions and response of local health authorities. Community case definitions are designed to align with the local language and do not require medical training to report on.
- Information discovered during surveillance should be shared with the local branch and health authorities based on the agreed protocol. Where relevant, (e.g. for zoonoses or environmental health events) information should also be shared with animal health and environmental health authorities.
- Community-based surveillance (CBS) can be done alongside other health, WASH or community engagement activities in your community, and therefore is not a stand-alone activity, but one that is useful to partner with other community-based activities.
- Community-based surveillance aids in
  - Early detection of public health risks within the community
  - Complementing early warning systems, extending them to the community
  - Linking early detection to early action within the community

## What to do and how to do it

- Preparation activities
  - Work with supervisors in mapping community needs and human, animal and environmental disease priorities (see Action tool [Community mapping](#))
  - Familiarize yourself with the disease that may be present in your community including signs and symptoms
  - Establish who is vulnerable in the community. Doing this will help you to identify people who are more likely to fall sick
  - Ensure referral mechanisms are clear in case community members fall sick and require referrals to health facilities for care.
  - Engage in community engagement activities such as mobile cinema, house-to-house visits, etc. to remain active and a known resource in the community.
- Recognize
  - Detect signs and symptoms corresponding to human, animal or environmental health risks or events in your community aligned with community case definitions
  - When you detect people who are sick with the disease, assess how severely ill they are and whether they need to be referred to a health facility (see Action tool [Referral to health facilities](#)).
  - Record the health risk or event you detected to ensure it can be followed-up

- Report
  - Report on the detected health risks or event in your community to your supervisor based on the methodology you are trained on (for example, SMS, phone call, or mobile application). *Remember that reporting must be systematic. To avoid confusion, everyone who reports should follow the same methods agreed on in the protocol and in the training.*
  - Your supervisor will then cross-check the report ensuring it meets the community case definition or unusual event requirements agreed on with health authorities. If matching, the supervisor will escalate the alert to the local health authorities for a response or investigation
  - After verification, the supervisor will notify relevant authorities in animal and environmental health for significant animal, zoonotic and environmental health events, especially those that portend a risk to human health.
- React
  - Begin community-level activities based on the health risk following proper safety precautions
  - Referral or care at home
    - Communicate specific health messages and information, and refer sick people promptly to health facilities
    - If sick people can be cared for at home, show their families what to do and provide them with information and supplies, where possible. Use corresponding “volunteer actions” in the ECV toolkit corresponding to the suspected epidemic risk.
  - Support health authorities in their investigation or response following-up on the alert
  - Where relevant, collaborate with and support officials in the animal and environmental health sectors for joint investigation, response and information sharing.

Additional resources on community-based surveillance: <https://cbs.ifrc.org/>

## Community messages



24. Finding sick people

## 02. Community mapping

### Overview

A map of the community enables you to connect issues or problems with particular places and makes information easy to see. Maps are often easier to understand than words.

Mapping aids in:

- Identifying risks and exposure to risk
  - Who and what are most exposed
  - What are they exposed to
- Show existing problems and vulnerabilities (some might make the current threat more serious)
- Understanding resources within the community that might be useful in managing the epidemic
- Obtaining information about other sectors (such as livelihoods, shelter, WASH, infrastructure etc.) that might be influenced by the epidemic, or that might be useful in managing the epidemic
- Analysing links and patterns in the exposure and spread of the epidemic - which may include human-to-human transmission dynamics, exposure from animals, vectors or food, behavioural risks, and environmental health drivers.

It is important to create the map together with community members. This helps communities to be active and to be participating members in the care offered by the Red Cross Red Crescent and volunteers.

Community mapping is especially useful in epidemics because it helps to see where the biggest problems and needs are and helps to identify risks and resources such as health posts, emergency vehicles, access roads, shelters, water sources, and so on. Maps can be used to support prevention, preparedness and response to an epidemic.

### How to make a community map

If possible, obtain or create a digital map of the community. If not available, it is possible to draw a simple spatial map that shows the community and all its key reference points. While keeping the fundamental principles of data protection, a community map should include the following:


- The whole community: concentrations of people, their houses, and who lives where
- The main shared/public locations in the community, such as schools, health centres, places of worship, water sources, markets, playgrounds and community meeting centres, communal livestock farming and livelihood areas such as cattle kraal, live bird markets, abattoirs, etc.
- The location of people who are most at risk [if you can identify them]
- Where the epidemic started and how it is spreading [if known and possible to identify]
- Health hazards and risks (e.g. improper rubbish disposal sites, large vector breeding sites)



## Using the community map

The map can be used to mark new cases and/or referred cases. Do as follows:

- Form teams to cover certain areas of the map.
  - Ensuring the participation of members from the community, each team should find out what it can about its area (how many people are sick, who is vulnerable, how many have been referred to health authorities, any other relevant information). If a zoonotic disease outbreak is suspected or implicated, find out who among the community members is keeping animals, how many animals are sick or have died, and/or presence of vectors in households, environment or even in surrounding water bodies. Work with your manager to target and prioritize those who are most at risk. This will require targeting geographically and, within those identified areas, targeting the most in need based on a vulnerability and capacity analysis that includes a gender and diversity analysis.
- Combine the maps of different teams. In doing this, you will be able to see:
  - Which areas of the epidemic you are covering, which areas you may not be covering, and details of each area. This will help you plan your actions. Some of these actions might include: environmental clean-up; distribution of bed nets; immunization campaigns; quarantine, animal biosecurity measures and other activities associated with managing the epidemic.



Making a community map

## 03. Communicating with the community

### Overview

Communicating during an epidemic can be difficult. Disease outbreaks, especially new ones, can cause uncertainty, fear and anxiety that can result in circulation of rumours, disinformation and misinformation. People may not trust the authorities, the health system or organizations including the Red Cross Red Crescent. They may not listen or may not believe the information they receive from people or organizations they do not trust. People may also be overcome with grief for those who are sick or who have died.

Sometimes, communities have strong beliefs that are different from the preventive and protective social measures promoted by the authorities and healthcare providers. They may believe strongly in their own cultural practices, traditional medicine, or other methods that might not prove effective against the disease. They may not accept certain treatments (including medicines and vaccines).

In many countries messages take the form of directives and one-way-communication. However, community engagement and participation have played a critical role in successful disease control and elimination campaigns in many countries

During a disease outbreak, trusted communication with the community is vital. To build trust, two-way communication is important. "Two-way" means volunteers should both *give messages to* AND *receive messages from* the community. Community members must feel respected and listened to and should have the opportunity to share their beliefs, fears and concerns. To accept volunteers' messages, community members must be able to trust you and have confidence in what you say. Once you understand the beliefs, fears and concerns of community members, you can provide them with truthful and accurate messages.

Providing health messages that are consistent, clear and easy to understand also helps to build trust. Giving accurate information to the community is critical, especially when it is necessary to persuade people to adopt safe practices (which might be different from what they would normally do). Some changes in behaviour that may be promoted are things such as:

- Accepting vaccinations or other medical treatments
- Washing hands with soap at crucial times
- Wearing personal protective equipment
- Burying loved ones in ways that are different from what they would normally do (safe and dignified burials)
- Practising social distancing
- Wearing insect repellent or sleeping under bed nets
- Agreeing to be isolated from others to avoid infecting them
- Preparing food and water differently (often by cleaning, boiling or cooking thoroughly)
- Quarantine and culling of animals (which in the case of livestock animals, is often a main source of food, nutrition and livelihood and may be difficult to accept by the farmers who own them)
- And other recommended public health measures

## What to do and how to do it

### *Communicating in an epidemic*

- Engage and involve community leaders and community members
  - Find out where the community obtains its information: Who do they trust to give them health information (for example: health authorities, community or religious leaders, doctors, traditional healers)
  - Work with communities to identify, choose and plan appropriate solutions for stopping the spread of disease
  - Talk to members of the community about their ideas, fears, beliefs and actions
    - Try to understand how much they know about the disease and its transmission
    - Try to understand beliefs and practices that might impact the spread of the epidemic
    - Try to understand what motivates or helps them to change behaviours
    - Try to understand what stops them from changing their behaviour
- Use different methods of communication
  - Use two-way communication when possible
    - When you understand the community's beliefs, fears and concerns, try to address these in your own messages
  - Sometimes, one-way communication methods are used to spread health messages to large numbers of people quickly
    - One-way communication methods should always be accompanied by two-way communication methods to ensure the community perspectives are known and listened to
  - People learn and retain information differently. It is important to use different methods
    - Communities are composed of different people and groups who may have different communication preferences or needs.
      - Think about how to target different groups, especially those who are hidden, stigmatized or considered "different" because of their religion, sexual orientation, age, disability, illness, or for any other reason:
        - Think about where you will go to reach them
        - Find out if they trust the same or different sources than other groups within the community
        - Discover if they have different access needs, such as language translation or in case of a disability – a different method of communication
    - When choosing methods of communication, consider what people prefer, trust and can access easily
      - Think about the characteristics of your target groups (for example, do they have access to media, such as radio or television? Can they read if they receive pamphlets of information and in what language? Are they accustomed to getting information from social media? Etc.)
      - Think about the resources you have access to (for example: do you have access to poster printing? Is there an appropriate location within the community where you can offer to answer questions or give out information? Etc.)
      - Consider the content of your message(s) and think about the most appropriate way to share that content in the specific context (for example: targeting men and women separately)
- Communication should be:
  - **Simple and short.** People should be able to understand messages easily and be able to remember and repeat them accurately and without difficulty.
  - **Trusted.** Delivered by people the community trusts, by a method the community trusts (for example:

radio, television, posters, town-hall discussions, market meetings etc.).

- **Accurate and specific.** Always provide correct and precise information. Messages should be consistent and should not be cause for confusion. *If* messages must change (due to new and advancing information about the epidemic), be honest and clear about what has changed and why.
- **Focused on action.** Messages should be action-oriented and should advise members of the community about what they can do to protect themselves and others.
- **Feasible and realistic.** Make sure that people have the capacity and resources to carry out the actionable advice you give.
- **Context-specific.** Information should reflect the needs and situation of the specific community. In all your messages, take account of social and cultural factors that might encourage community members to adopt safer behaviours (such as accepting vaccines) or prevent them from doing so.

### ***Different ways of communicating***

There are many, many ways to communicate with communities. The following one and two-way methods of communication are some examples you might consider. Methods can (and should) be combined to ensure accessibility to as many community members as possible.

- One-way communication methods
  - Video, films, television commercials
  - Songs, poems, drama, role-play or theatre, or other edutainment methods
  - Community announcements such as: community town-criers, loud-speaker announcements, SMS or WhatsApp, mass messaging, social media messages, radio broadcasts
  - Posters, billboards
- Two-way communication methods
- Door-to-door visits
- Meeting with key informants such as: community or religious leaders; traditional healers or midwives; teachers; elders, etc.
- Community discussions encouraging participatory methods such as: three pile sorting, voting charts, mapping, polling, barrier analysis, community planning
- Use of feedback and suggestion boxes or presence of trusted focal persons to receive anonymous feedback or messages from community members.

### ***Pay attention to rumours***

Rumours can cause panic and fear or can promote unsafe practices. Under the influence of the rumours, communities can lose trust in the health authorities, and they may lose belief in the ability to stop the epidemic. Rumours sometimes cause people to reject interventions that could prevent the spread of disease. Volunteers must:

- Listen for rumours or incorrect information.
  - Note when and where a rumour was heard and report it to your volunteer supervisor or National Society focal point immediately
  - Try to understand why the rumor is spreading fast and of what importance it is to the community. For example, is it just based on lack of knowledge or fear of the unknown, or is it associated with certain socio-cultural beliefs or associated with the stigmatization of a certain demography of people?
- Correct the rumour

- Give the community clear, simple facts about the disease
- Reiterate and explain clearly what they can do to protect themselves and others

## 05. Volunteer protection and safety

### Overview

Volunteers work in vulnerable situations and with people of many capacities. Working in epidemics can be risky because volunteers can also catch a disease and fall sick. In addition to physical risks, there may be risks to volunteers' emotional and mental well-being, due to the nature of the work they undertake. It is important to protect from and minimize the impacts of these risks.

Your National Society should provide proper protection for you and other volunteers who are working in epidemics. Your manager is a valuable resource for information and equipment to protect and preserve your physical, emotional and psychosocial wellbeing.

It is important to follow the guidance from your supervisor and National Society and use the level of protection that is appropriate for the situation you are in.

### What to do and how to do it

#### *Protecting yourself and others from disease*

1. You must be familiar with and trained to use protective equipment before you wear it in an actual disease environment. Try the equipment out beforehand and learn how to use it properly.
  - In certain epidemics like Ebola, Marburg, Lassa fever and plague, full protection should be used whenever you undertake high risk activities. Full protection requires use of personal protective equipment (PPE). (See Action tool [Personal protection equipment \(PPE\) for highly infectious diseases](#))
  - In other epidemics, you should at least use masks and latex gloves and wash hands with soap after contact with an affected person or animal. (See Action tool [Handwashing with soap](#) for instructions in good hand hygiene.)
2. Volunteers should be vaccinated according to country-specific vaccination guidelines (see Action tool [Routine vaccinations](#)).
  - Volunteers should be vaccinated according to the routine vaccination schedule in the country
  - Volunteers may be eligible for vaccination during mass vaccination if applicable
3. Volunteers should be alert to their own physical and psychosocial well-being during an epidemic
  - Volunteers should be alert to stressors in their personal and working lives, and should have a plan in place for how to cope with stress and trauma in a healthy and helpful way
    - This may include stress management techniques that you already use such as exercise, meditation, taking part in hobbies, etc.
    - Your manager is a useful resource for information and tools to use to help you achieve and maintain psychosocial well-being

#### *Understanding common prevention and control measures*

Volunteers should learn additional prevention measures for use in epidemics (and before them). These include:

- Vector control measures (see Action tool [Vector control](#))
- Safe handling of animals (Action tool [Handling and slaughtering animals](#))
- Chemoprophylaxis (Action tool [Chemoprophylaxis](#))
- Safe food and water (Action tools [Good food hygiene](#) and [Clean, safe household water](#))
- Hand hygiene (Action tools [Handwashing with soap](#) and [Handwashing in a highly infectious epidemic](#))

## Protecting volunteers from harm and liability to others

1. **Volunteers should be protected** if they suffer damage or injury in the course of their work. Accidents can happen, and volunteers can be injured or even killed. Equally, volunteers can harm other people and their property, especially if they have not been properly trained or given the correct equipment.
  - National Societies therefore need to have appropriate insurance policies. Insurance may be needed to pay compensation to volunteers or their families if they are injured or killed; to pay compensation to others if they suffer harm as a result of volunteer actions; and to cover legal costs. The nature of the cover will depend on the legal system in your country. The Movement urges the National Societies to acknowledge and uphold their duty of care towards volunteers, especially if something should happen to them while carrying out their duties. Ask your manager about the type of insurance or coverage through a “safety net” you can receive.
  - Prior to asking volunteers to perform activities that are high-risk (e.g. safe and dignified burials), National Societies should also ensure volunteers have received the necessary vaccinations and protective equipment. What this includes will depend on the context in which you are working and the health policies for staff and volunteers of your National Society.
2. **Volunteers should be informed** of and understand the National Society's security policy and follow the rules and regulations it sets out. You should also be informed of any changes in the policy and asked to report any incidents of concern.
  - Safety in the community depends on the personal attributes of volunteers, trainers and other team members – how they work together and how they collaborate with people in the community. Volunteers should be culturally sensitive. Your personal behaviour should never cause offence. You should show integrity and should never become a problem for the community. Correct, polite, impartial behaviour is always expected.
  - Volunteers should be proactive in managing and maintaining their own safety and security. This means you should not hesitate to ask your manager about safety and security risks and what you should do if you encounter any threats or have any problems. You should find out what protocols are in place if a safety or security incident occurs, including how and to whom you should report these events.







## 06. Personal protection equipment (PPE) for highly infectious diseases

### Overview

When working with certain epidemic diseases, especially highly infectious diseases such as Ebola, Marburg, Lassa fever and plague, it is vital to protect yourself when you touch affected people, their body fluids, or dead bodies or animals.

Like volunteers, PPE comes in different sizes and fits. Not every size of PPE will fit every volunteer appropriately (e.g. protective goggles may not be the right size or shape for some women or for people of certain ethnicities, leaving them at greater risk of exposure). It is very important that volunteers have the correct size and fit of all their equipment for optimum protection.


This tool shows you how to put on and take off PPE. \*\* Note: Different PPE types are used in different contexts, depending on the disease and the type of protection required (e.g. the components of the PPE for Ebola and for plague may have some commonalities but are not the same). Always ensure that you have been properly trained to use the type of PPE available in your context, for the specific disease you are addressing. \*\*

See Action tool [Volunteer protection and safety](#) for more information on volunteer protection.


### What to do and how to do it

- Test the size and fit of your PPE by having your manager check the fit and coverage of all your equipment. If anything is incorrectly sized, be sure you obtain equipment of the correct fit before you begin work that puts you at risk of exposure.

### Steps for putting on protective clothing



**Steps for taking off protective clothing**



**Note:** Handwashing, with gloved hands, should be performed before starting to remove protective clothing (step 4), between removing protective clothing but before removing protective face mask (step 9) and at the end with ungloved hands (step 12).

## 20. Isolating sick people

### Overview

- Some diseases are so dangerous or infectious that it is necessary to isolate people who are sick to prevent them from passing the infection to others. Isolation means separating those who are sick from those who are healthy.
- Isolating people is not imprisonment; it is not done forcibly and should be carried out with their consent. You need to explain to affected people and their families why isolation is necessary. People may also need to be isolated from their companion or livestock animal (and vice versa), if it brings the risk of transmission of disease. In extreme cases of progressive outbreaks, sick livestock animals may need to be culled (selectively slaughtered) to stop the spread of a disease to other animals and humans. Animal isolation, quarantine or culling should also not be done forcibly and should be carried out with their care-giver's consent.
- When a person is in isolation, the people who come into direct contact with them (for example, family members and healthcare providers) must be protected using the appropriate protective measures, like personal protective equipment (PPE). The number of people who are in direct contact with the sick person should be kept to a minimum. For example, only one family member should provide care, food, water, etc. to the sick person.

### What to do and how to do it

#### *Surveillance and detection*

1. Even if a disease requires isolation, you should continue monitoring, looking for cases of the disease (see Action Tool [Community-based surveillance](#)).
  - When cases are found, the need for isolation should be explained to those who are ill and their families.
  - Explaining isolation protects others, including members of the sick person's family. It may be easier for both the sick person and his or her family to understand why isolation is being proposed if they receive a complete explanation.
2. Provide support to the affected person, family members and healthcare providers to facilitate isolation.
  - This is not done forcibly, and the wishes of the family and sick person should be respected as much as possible.
  - Those who care for a person in isolation should be shown how to assist him or her and how to protect themselves. They should be provided with proper protective equipment.

#### *Management and care*

1. Provide psychosocial support to families (see Action Tool [Psychosocial support](#)). Isolation of a loved one can be very hard, for the family as well as the person in isolation.
2. Make sure that people in isolation have adequate food, water, and health care, as well as help to cope with any loss of livelihood.
3. Ask if those who need isolation have dependants (e.g. children, people with disabilities) that may need additional support. Inform your manager, who will liaise with the Red Cross Red Crescent protection

team for dependants to get the appropriate support.

### ***Social mobilization, messaging, and community engagement***

1. Talk to the family, community and elders/leaders to prevent stigma and social rejection.
  - Try to stop isolation from causing stigma or social rejection of sick people and their families.
  - Explain to the community why isolation is necessary.
  - Seek the help of elders and community leaders to fight stigma.



Talk to the community and elders to prevent stigma and social rejection.

## **Community messages**



24. Finding sick people

## 26. Respiratory hygiene and coughing etiquette

### Overview

- Many diseases are spread through particles or droplets that get in the air when people cough, sneeze, blow their noses or spit. There are some easy ways to minimize the spread of these diseases by practising good respiratory hygiene and coughing etiquette.
- The correct way to cough or sneeze without transmitting germs is to use a disposable tissue, a handkerchief or your sleeve. When you cough or sneeze into these, you are less likely to spread germs. If you use a disposable tissue, dispose of it safely as soon as possible. If you use a handkerchief, be sure to wash it frequently. Always wash your hands as soon as possible if you use a handkerchief or tissue when you cough or sneeze.
- Many people cough or sneeze into their hands to stop germs spreading. This is not the best practice. Afterwards, you can transmit the germs by touching things or shaking the hands of other people. The best thing to do is avoid coughing or sneezing into your hand. If you must use your hands, wash your hands with soap and water as soon as possible and without touching anything.
- It is also important to blow your nose and/or spit in a safe way. Tissues or handkerchiefs should always be used when you blow your nose or spit. Do not practise open spitting (for example, spitting onto the ground or into spittoons), always use a tissue or handkerchief (tissues are preferable and should be disposed of safely as soon as they are used).

### What to do and how to do it

#### *Hygiene promotion*

Promote respiratory hygiene and coughing etiquette to people in your community.

- Ask your schools and other places where people gather in groups if they would like you to come and speak about respiratory hygiene and coughing etiquette.
- Explain that coughing etiquette matters because it helps to prevent the transmission of diseases that are spread through droplets carried in the air.
- Show people how to cough properly and ask them to teach the same behaviour to others.
  - When talking about coughing etiquette, also teach people about proper hygiene and social distancing (see Action Tools [Social distancing](#) and [Handwashing with soap](#)).
  - Make posters that show the DOs and DON'Ts of coughing etiquette and put them up around your community. (See Action Tool *Social mobilization and behaviour change* for more information on social mobilization techniques).
- In some countries, chewing Khat or chewing tobacco is widely practised. In these places, volunteers should explain that gathering in enclosed and poorly ventilated areas to chew and spit (khat or tobacco) for extended periods of time may increase the spread of germs and increase the risk of transmission of respiratory infections.





## Community messages



14. Vaccinations for children





18. Coughing correctly

## 27. Shelter and ventilation

### Overview

The environment and spaces people spend time in have a large impact on their health and well-being. Many diseases can spread through air or because of water and sanitation conditions.

- Some diseases are spread by droplets in the air (often through coughing and sneezing). Germs can become more concentrated in poorly ventilated rooms or homes in which a sick person is coughing or sneezing. In buildings that are stuffy and poorly ventilated (with little flow of air) a sick person can infect every other person in the same space. To reduce the risk of spreading disease to others, it is a good idea to ventilate houses. People should be instructed to open windows or doors to allow fresh air to come in and stuffy, old air to go out.
- When many people live together in the same space (overcrowding), they can also catch infections more easily from one another. If possible, people should have enough space to move and breathe freely. If possible, encourage people to live or gather in places where there is adequate space.
- In addition to the risk of uncontrolled fires and burns, cooking with charcoal or firewood in an enclosed space (a room or shelter without windows) can harm people who are sick with air-borne diseases. Smoke can make it difficult to breathe, especially if other respiratory conditions are present. It is important to have enough air flow to take the smoke and fumes away.
- Other diseases are spread via contaminated water or bad sanitation or absence of shelter. Diseases that spread via contaminated water or bad sanitation (diarrhoeal diseases, cholera, typhoid, hepatitis E, for example) will spread more easily if clean water is lacking or if sound, clean toilets are not set close to where people shelter. Proper shelters, with clean water and sanitation facilities, which protect people from rain, wind, sun and cold help prevent diseases from spreading and help sick people recover from disease.

### What to do and how to do it

#### *Preventing disease in shelters*

- When people move following conflict or disaster, the shelters they move into are not usually as good as those they are used to. Help people to have the best possible shelter, one that is well ventilated and has a good water supply, with latrines and waste disposal facilities close by.
- Always ventilate shared or communal shelters and emergency accommodation.
- Advise people to keep their windows open if the weather permits, especially if they are caring for a sick person.
- Encourage people to wash their hands after going to the toilet, after cleaning a child, and before preparing or eating food.
- Encourage people to wash water storage containers regularly with soap and water, and to keep the area around the shelter free from rubbish and animal waste.

#### *Managing disease in shelters*

- When an outbreak of air-borne diseases occurs, it is important to tell your community about the importance

of good ventilation and encourage them to open windows and doors regularly.

- When a food or water-borne disease outbreak occurs, it is important to tell your community about the importance of food hygiene and safety, as well as the importance of access to clean water and proper sanitation facilities and waste disposal.

### ***Social mobilization, messaging and community engagement***

- Talk to the community about the importance of a clean water supply, sanitation, handwashing and good food hygiene. (See Action tools [\*Clean, safe household water\*](#), [\*Good food hygiene\*](#), [\*Sanitation\*](#), [\*Building and maintaining latrines\*](#), [\*Handwashing with soap\*](#) and [\*Handwashing in a highly infectious epidemic\*](#)).



## **Community messages**



22. Good ventilation

## 29. Hygiene promotion

### Overview

Hygiene promotion is a term used to cover a range of strategies aimed to improve people's hygiene behaviour and prevent the spread of disease. Hygiene promotion enables people to take action to prevent water, sanitation and hygiene-related diseases by mobilizing and engaging the population, their knowledge, and resources.

The focus of hygiene promotion is determined based on the health risks. By creating a series of barriers to infection, hygiene behaviour has a critical influence on the transmission of water- and sanitation-related diseases as shown in the 'f' diagram<sup>[1]</sup> below:

# The 'F' Diagram


The movement of pathogens from the **faeces** of a sick person to where they are ingested by somebody else can take many pathways, some direct and some indirect. This diagram illustrates the main pathways. They are easily memorized as they all begin with the letter 'f': **fluids** (drinking water) **food**, **flies**, **fields** (crops and soil), **floods**, **fingers** and **faces** (and surface water generally).

★ **WATER**

★ **SANITATION**

★ **HYGIENE**

Barriers can stop the transmission of disease; these can be primary (preventing the initial contact with the faeces) or secondary (preventing it being ingested by a new person). They can be controlled by water, sanitation and hygiene interventions.



**Note:** The diagram is a summary of pathways; other associated routes may be important. Drinking water may be contaminated by a dirty water container, for example, or food may be infected by dirty cooking utensils.

© WEDC

**Source:** McMahon, Glenda; Davey, Kay; Shaw, Rod (2020): P004 The F Diagram. Loughborough University. Poster. <https://doi.org/10.17028/rd.lboro.12738692.v1>

## What do to and how to do it

### ***Understand the community***

1. Familiarize yourself with the ways in which people collect water, store food and water, dispose of rubbish, wash themselves and use latrines.
  - Identify key places where hygiene is important not only on an individual, but also on a communal level, such as markets, schools, restaurants or churches. Find out if you can work with them to promote good hygiene practices.
    - You might hold a hygiene session for students or teachers or help the market sellers to build a handwashing station and outside latrine for users.
2. Have a conversation with members of your community about hygiene.
  - Include women, community leaders, caregivers and decision-makers.
    - Make sure they understand that good hygiene is important and can stop the spread of disease.
3. Be a role model for others in your community. Use a clean latrine, dispose of your rubbish, wash your hands often.

### ***Promote community hygiene messages***

Normally the key issues to address include the following. Click on the corresponding action cards to obtain the information you need:

- [Food hygiene](#)
- [Clean and safe household water](#)
- [Personal and hand hygiene](#)
- [Environmental sanitation](#)
- [The control of flies, mosquitoes and other disease vectors](#)

---

[1] Shaw, R. 2013. *The 'f' diagram - Landscape*. WEDC Graphics: Disease, Water, Engineering and Development Centre (WEDC): Loughborough University, UK.

## Community messages



04. Storing water properly



05. Using clean safe drinking water




06. Using a clean latrine



08. Washing hands with soap





09. When to wash hands



11. Cleaning up places where mosquitoes breed



12. Good food hygiene



13. Good personal hygiene



20. Collecting and disposing of rubbish

## 34. Handwashing with soap

### Overview

Handwashing is one of the most important ways to prevent the spread of several epidemics, especially diarrhoeal diseases. Handwashing is easy and everyone (including children) can and should do it. To wash hands people must have access to water and soap.

Hands should be washed with soap:

- **BEFORE:**
  - Preparing food
  - Eating
  - Feeding a child
  - Breastfeeding
  - Caring for someone who is ill or treating a wound (yours or someone else's)
- **AFTER:**
  - Using the toilet
    - Men, boys, women and girls should wash their hands after using the toilet
    - Women and girls should engage in menstrual hygiene during their monthly menstrual cycles
      - Promote the use of clean, dry materials (disposable or reusable)
      - Promote changing menstrual materials and bathing as often as needed.
      - Discourage sharing reusable pads with anyone else
  - Cleaning a baby
  - Touching garbage or waste
  - Touching or feeding animals; handling raw meat
  - Blowing nose, coughing or sneezing
  - Treating wounds or caring for sick people
  - Coming into contact with a sick person in an epidemic (see Action Tool *Handwashing in a highly infectious epidemic*)

### What to do and how to do it

#### ***How to wash hands***

1. Wet your hands and rub them with soap.
2. Rub all parts of your hands together for 10 to 15 seconds.
3. Use lots of force (push your hands together hard) and remember to wash all surfaces, including the backs of the hands and between the fingers.


4. Rinse hands well so they are free of soap.
5. Dry hands with a paper towel. If there is no towel, wave them in the air until they are dry.



## Community messages



08. Washing hands with soap



09. When to wash hands

## 35. Handwashing in a highly infectious epidemic

### Overview

- Handwashing is a vital source of protection in epidemics of highly infectious diseases such as Ebola, Marburg fever, Lassa fever, plague, MERS and monkeypox. It is possible to become exposed to these diseases and to fall sick when trying to help people. Highly infectious germs are spread very easily through body fluids and certain diseases can even be transmitted via the bodies of people who are dead.
- Very vigorous and comprehensive handwashing is needed to protect yourself. (See Action Tools: [\*Volunteer protection and safety\*](#); and [\*Personal protective equipment \(PPE\) for highly infectious diseases\*](#))

### What to do and how to do it

#### ***When to wash hands in a highly infectious situation***


Hands should be washed with soap and/or disinfectant:

- BEFORE: preparing food, eating, feeding a child, breastfeeding, caring for someone who is ill, treating a wound (yours or someone else's)
- AFTER: using the toilet or cleaning a baby; touching garbage or waste; touching or feeding animals; handling raw meat; blowing nose, coughing or sneezing; treating wounds or caring for sick people; coming into contact with a sick person in an epidemic
- IN ADDITION: During an epidemic (of a highly infectious disease, or another disease such as diarrhoea or cholera, or a respiratory infection), it is a good idea to use this handwashing method every time you have touched something that can pass on an infection.

#### ***How to wash hands in highly infectious epidemics***

When working in highly infectious epidemics, it is critical to use all available forms of protection. One of these is handwashing. To wash hands correctly during a highly infectious epidemic, you must:

- Use soap or an alcohol-based disinfectant.
- Wash your hands as usual.
- Then wash thoroughly between your fingers.
- Scrub the tips of the fingers of the two hands together.
- Wash each of your thumbs with the other hand.
- Scrub the tips of the fingers of each hand on the palm of the other hand.



## Community messages



10. Steps for washing hands in epidemics

## 43. Social mobilization and behaviour change

### Overview

There are many reasons why people practise unhealthy behaviours. People are affected by access to services or facilities, social norms and influences where they work, live or play. Behaviour change is the study of how and why people change some habit or action in their life. As volunteers, we need to understand WHY the behaviour is happening and WHAT actions will lead to change to create healthy behaviours. Examples of healthy behaviours include handwashing, breastfeeding, immunizations, consistent condom use and use of bed nets.


In any culture and context, behaviour change involves three elements. Before people will change their behaviour:

1. They need to know what, why and how they should change. They need knowledge.
2. They need to have the right equipment, access and capacity. They need an enabling environment.
3. They need to be motivated to change.





The social-ecological model below shows how each person's behaviours are affected by many different levels of influence including the individual level, the interpersonal level, the community level, the organizational level and the broader policy level which includes laws and policies that allow or restrict a behaviour. In order to promote health, it is important to consider and plan behaviour change activities across multiple levels at the same time. This approach is more likely to result in successful behaviour change over time. As a volunteer, it is helpful to understand that behaviour change is difficult for many people because of these many levels and the complex interactions and expectations across the different levels. If you consider how each of the levels affects the behaviour of the person you want to help, you can try different interventions at each level that is specific to their needs.



Socio-ecological model

### What do to and how to do it

The general process for developing a behaviour change intervention includes staff and volunteers working through the general steps of:

1. Sensitizing the community to the behaviour change process using the theory of change model.
2. Assessing the problem behaviour – why it is practised, who practises it, when it is practised and what factors in the environment or society encourage the behaviour. Assess this information at the different levels of the social-ecological model for each community you serve.
3. Identifying an appropriate behaviour goal based on your assessment.
4. Reviewing the causes or barriers at each level that allow the behaviour to continue. Identify interventions

that align with each cause or barrier and that can be used at different levels.

5. Discussing the suggested interventions for each social-ecological model level with the community.
6. Identifying appropriate interventions for the context at each level. Interventions should be planned to address the stages of the theory of change by first giving knowledge and addressing environmental factors, motivating key people to gain approval and intentions, and ultimately inciting people to action that contributes to the overall goal.
7. Implementing the interventions at each level.
8. Monitoring to see if change is happening. Change takes time but it must be monitored to ensure that it is happening, even slowly. Additionally, as people go through the change process, their barriers and causes will change. The behaviour change interventions should adjust to these changes to ensure that change can continue.
9. Recognizing that when change is not happening as intended, further assessment and intervention tweaking is needed.
10. Continuing to implement, monitor, assess and adjust as the change process happens.

For more information, please consult the eCBHFA Manual for volunteers on [Behaviour Change](#), including:

1. Principles of behaviour change
2. The social ecological model
3. The stages of behaviour change
4. Activities for behaviour change

## Community messages



23. Encouraging healthy behaviours in a community

## 44. Dealing with rumors

### Overview

- Rumours are stories of uncertain truth. They often spread in communities during epidemics when people feel fear or anxiety about the disease. Rumours often provide an explanation for what is unknown about the disease, even if the explanation is not true.
- Rumours include *misinformation* or *disinformation*. Misinformation is when incorrect information is spread, without the intent to deceive, through a misunderstanding or a mistake. An example of misinformation is the rumour that Ebola is caused by witchcraft. In truth, Ebola is spread by a virus, but people often mistake it for witchcraft because they cannot see the virus or have never heard of the Ebola virus before.
- Disinformation is when incorrect information is spread intentionally to deceive or manipulate, such as “fake news”, which is disinformation disguised as news and is often spread for political or economic gain. An example of disinformation is when someone who sells multivitamins advertises that the tablets “cure” HIV, even though they know it is not true.

In epidemics, we often see two kinds of rumours:

- Rumours about possible cases
  - These rumours can violate community members’ privacy and right to confidentiality and may put them at risk
    - Rumours can often reflect pre-existing fears and prejudices within the community. This may lead to placing blame on different people or groups. This type of untrue belief can give community members “permission” to discriminate against someone or a group without feeling guilty, because of untrue beliefs.
  - They may also cause the unnecessary use/waste of health resources when rumoured cases must be followed up
- Rumours about the causes or treatment of the disease
  - Can distract from public health messaging
  - May conflict with the behaviours and practices recommended to fight the epidemic
  - Can create a dangerous situation for volunteers and healthcare providers if they cause mistrust

Paying attention to rumours can help us to understand the beliefs and perceptions that influence people. Using this information, we can make our messages specific to the community, the context and the beliefs. Rumours may also serve as a warning sign of hazards such as violence or risky behaviours so these can be managed quickly.

### What to do and how to do it

#### *Listen for rumours and capture information*

- Establish a system for listening to rumours: Listening for rumours involves more than just hearing the words people use. To effectively listen to rumours, you need to:
  - Build trust with community members: Identifying rumours is not as simple as asking people about any

rumours they have heard. This will not necessarily uncover rumours because people may believe a rumour to be true and therefore not consider it a rumour. Also, people may not trust you as someone to discuss their beliefs with, in this way.

- Listen to the language the community is most comfortable using
- Tune into social and traditional media to understand what people are hearing and what they are saying
- Engage in open and unstructured conversations with diverse groups of people to understand the beliefs they hold and why they hold them
- Host group discussions with community members and members of community groups (such as women's or youth groups)
- Pay attention to what you hear during your work as a volunteer, but also during your personal time
- Establish a method of collecting information about rumours: Use a rumour log where you can record:
  - Details – what is the rumour?
  - Date - when was the rumour heard?
  - Location - where was the rumour heard?
  - Channel – how is the rumour being shared/spread?

### ***Report and help to verify rumours***

- Report rumours to your supervisor: Ask them to verify if the rumour is true or false
  - Sometimes parts of a rumour are true and other parts are false. It is important to understand the facts
  - Follow the guidance of your supervisor in uncovering more information about the rumour if possible
  - You may be asked to find out more information about the rumour from community members. You may also be asked to speak to the source of the rumour to understand more about what is being said and why
    - Explain that you are verifying a rumour, which may or may not be true, and repeat the rumour that you have heard
    - Ask them what is true/untrue about the rumour you heard and to state in simple terms the facts and how they know them
    - Repeat what you have heard, to check that you have understood them correctly. You should finish with a clear understanding of what they are describing – if you are not sure ask again
    - Try to find out what triggered the rumour. For example, did the rumour start because of a badly worded message? A government announcement? Etc.

### ***Plan a response to rumours***

- With your supervisor, develop a plan to address and prevent rumours. Do not ignore or deny rumours.
  - Rumours usually do not go away on their own and can cause big problems if they are not addressed
  - Replace rumours with accurate information
    - Respect local customs and beliefs and align messages with pre-existing beliefs and customs. For example, a common rumour might be that Ebola is caused by witchcraft; the conventional response is to refer to Ebola as a virus. However, rather than deny this pre-existing belief, it may be more useful to accept that this is what people believe and to create recommendations and messages that align such as: do not touch this person unprotected but feel free to provide food [and prayers] as a token of empathy
    - Use communication channels/people that community members trust
    - Use language that people understand and are comfortable with

- Continue to engage in conversation with communities to make sure you are being understood