



Zika virus infection

Last update: 2023-06-23

Key facts

Transmission: Vector-borne (mosquito)

- Mosquito bite (mostly spread by daytime biting mosquitoes, especially in early morning and late afternoon or evening)
- Mother-to-child transmission during pregnancy
- Unprotected sex with a person infected with Zika virus infection

Most vulnerable to severe consequences

- Pregnant women, women and girls of reproductive age, and couples who want children must remain very cautious - getting Zika during pregnancy can cause serious brain defects in an unborn baby

Most vulnerable to contracting the disease

- Any person in the community living or working near mosquito-breeding sites (stagnant water) can get Zika

Symptoms

(** People show symptoms with different levels of severity. Although some people become very sick, most people do not. Some people develop mild symptoms and some people do not show any symptoms, but they can still transmit the disease. Prevention of transmission is very important. **)

- Fever
- Skin rash
- Eye infection (red, draining eyes)
- Muscle and joint pain
- Feeling unwell
- Headache
- Weakness

What can you do to prevent and control an epidemic?

Vector control and prevention

- Initiate elimination of mosquitos and breeding sites
 - Remove standing water and apply larvicides
 - Promote community clean-up campaigns to remove rubbish and cover water containers
- Prevent mosquito bites by advocating the use of:
 - Insecticide-treated curtains or screens on windows and doors
 - Personal protection (application of repellents, wearing long sleeved clothes)
 - Insecticide-treated bed nets for children and others who sleep during the day

Monitoring the community and identifying sick people

- Identify community members with suspected Zika virus
- Identify babies born with microcephaly (small heads) or brain defects

Treatment and management

- Refer all suspected cases in pregnant women to health facilities
- Refer suspected cases of congenital Zika syndrome to health facilities
- Reproductive health counselling and support for couples who plan to have children and who live in areas where Zika is endemic
- Provide psychosocial support to the sick person and their family members
 - Promote ongoing psychosocial support to mothers, families and babies born with brain defects

Social mobilization and health promotion

- Find out the specific advice being given by health and other relevant authorities
 - Promote and practise public health messaging:
 - Encourage the practice of safe sex (especially applies to pregnant women, their partners, and travellers who return from areas in which Zika is present)
 - If possible, pregnant women should be encouraged not to travel to areas in which Zika is present
 - Identify if there are any community spaces where women give birth and engage with traditional birth attendants to share information about the disease transmission and prevention modes
- Model following this advice and inform community members of current health practice advice
- Offer support and encouragement to follow the advice
 - Try to gain understanding about if and why health practice advice is not being followed
 - With the guidance of your supervisor and health authorities, work with communities to overcome barriers to following health advice and recommended practices

Mapping and community assessment

- Make a map of the community.
- Mark the following information on the map:
 - How many people have fallen sick with Zika? Where?
 - How many babies have been born with microcephaly or other obvious brain defects? Where? When?
 - Who and where are the vulnerable people?
 - Where are the local health facilities and services? (include traditional healers)
 - Where do women give birth? (include traditional birth attendants)
- Record the following information on the back of the map:
 - When did people start to fall sick with Zika?
 - Have any children been born with brain or head defects?
 - How many?
 - How many people live in the affected community?
 - How many young women/pregnant women live in the area?
 - Have the authorities established a vector control programme?
 - Do people generally cover their water containers (inside and outside)? Who is responsible for the maintenance of containers for household drinking water and for vessels to do laundry; is it women or men?
 - How does the community usually remove standing, stagnant water?
 - What are the usual ways of disposing of rubbish and solid waste in the community?
 - How common is it for people to live in houses with insect screens on windows and doors?
 - How common is it for people who sleep during the day (such as babies and children) to sleep under insecticide-treated bed nets?
 - Do pregnant women usually sleep under a bed net?
 - Are nets hung up and maintained properly?
 - If people are not using nets for daytime sleep, why not?
 - What are the community's habits, practices and beliefs about repellents and sprays?
 - What are the community's habits, practices and beliefs about caring for and feeding sick people? When babies and infants are sick, do women continue to breastfeed?
 - Are children badly affected by Zika? Are there other groups (specific ages, occupations, geographic areas, etc.) that are badly affected?
 - Is a social mobilization or health promotion programme in place?
 - Which sources do people use/trust the most for information?
 - Are there rumours or misinformation about Zika? What are the rumours?
 - Are there family planning services that both women and men can access?
 - Who spends more time in the household during the day (and is more exposed to the mosquito bite)? Women, or men, or both?

Volunteer actions

- [01. Community-based surveillance](#)
- [02. Community mapping](#)

- [03. Communicating with the community](#)
- [04. Community referral to health facilities](#)
- [05. Volunteer protection and safety](#)
- [19. Mental Health and Psychosocial support \(MHPSS\)](#)
- [36. Vector and reservoir control](#)
- [37. Mosquito nets / bed nets](#)
- [38. Waste disposal and clean-up campaigns](#)
- [42. Promoting safe sex](#)
- [43. Social mobilization and behaviour change](#)

Other resources

Community-based surveillance and monitoring for Zika toolkit: **[Community based Surveillance and Monitoring for Zika Toolkit](#)**

Zika, Dengue and Chikungunya toolkit: [Zika, dengue and chikungunya toolkit | IFRC](#)

01. Community-based surveillance

Overview

- Community-based surveillance is the systematic detection and reporting of significant public health events (such as sudden illness or death in people or animals) within a community by community members and volunteers. It is a simple, adaptable, low-cost public health initiative designed to complement early warning systems for potential epidemic diseases.
- Volunteers use something called a “community-case definition” to detect and report signs and symptoms of potential diseases, health risks and events, and support in community actions and response of local health authorities. Community case definitions are designed to align with the local language and do not require medical training to report on.
- Information discovered during surveillance should be shared with the local branch and health authorities based on the agreed protocol. Where relevant, (e.g. for zoonoses or environmental health events) information should also be shared with animal health and environmental health authorities.
- Community-based surveillance (CBS) can be done alongside other health, WASH or community engagement activities in your community, and therefore is not a stand-alone activity, but one that is useful to partner with other community-based activities.
- Community-based surveillance aids in
 - Early detection of public health risks within the community
 - Complementing early warning systems, extending them to the community
 - Linking early detection to early action within the community

What to do and how to do it

- Preparation activities
 - Work with supervisors in mapping community needs and human, animal and environmental disease priorities (see Action tool [Community mapping](#))
 - Familiarize yourself with the disease that may be present in your community including signs and symptoms
 - Establish who is vulnerable in the community. Doing this will help you to identify people who are more likely to fall sick
 - Ensure referral mechanisms are clear in case community members fall sick and require referrals to health facilities for care.
 - Engage in community engagement activities such as mobile cinema, house-to-house visits, etc. to remain active and a known resource in the community.
- Recognize
 - Detect signs and symptoms corresponding to human, animal or environmental health risks or events in your community aligned with community case definitions
 - When you detect people who are sick with the disease, assess how severely ill they are and whether they need to be referred to a health facility (see Action tool [Referral to health facilities](#)).
 - Record the health risk or event you detected to ensure it can be followed-up
- Report
 - Report on the detected health risks or event in your community to your supervisor based on the methodology you are trained on (for example, SMS, phone call, or mobile application). *Remember that reporting must be systematic. To avoid confusion, everyone who reports should follow the same methods*

agreed on in the protocol and in the training.

- Your supervisor will then cross-check the report ensuring it meets the community case definition or unusual event requirements agreed on with health authorities. If matching, the supervisor will escalate the alert to the local health authorities for a response or investigation
- After verification, the supervisor will notify relevant authorities in animal and environmental health for significant animal, zoonotic and environmental health events, especially those that portend a risk to human health.
- React
 - Begin community-level activities based on the health risk following proper safety precautions
 - Referral or care at home
 - Communicate specific health messages and information, and refer sick people promptly to health facilities
 - If sick people can be cared for at home, show their families what to do and provide them with information and supplies, where possible. Use corresponding “volunteer actions” in the ECV toolkit corresponding to the suspected epidemic risk.
 - Support health authorities in their investigation or response following-up on the alert
 - Where relevant, collaborate with and support officials in the animal and environmental health sectors for joint investigation, response and information sharing.

Additional resources on community-based surveillance: <https://cbs.ifrc.org/>

Community messages



24. Finding sick people

02. Community mapping

Overview

A map of the community enables you to connect issues or problems with particular places and makes information easy to see. Maps are often easier to understand than words.

Mapping aids in:

- Identifying risks and exposure to risk
 - Who and what are most exposed
 - What are they exposed to
- Show existing problems and vulnerabilities (some might make the current threat more serious)
- Understanding resources within the community that might be useful in managing the epidemic
- Obtaining information about other sectors (such as livelihoods, shelter, WASH, infrastructure etc.) that might be influenced by the epidemic, or that might be useful in managing the epidemic
- Analysing links and patterns in the exposure and spread of the epidemic - which may include human-to-human transmission dynamics, exposure from animals, vectors or food, behavioural risks, and environmental health drivers.

It is important to create the map together with community members. This helps communities to be active and to be participating members in the care offered by the Red Cross Red Crescent and volunteers.

Community mapping is especially useful in epidemics because it helps to see where the biggest problems and needs are and helps to identify risks and resources such as health posts, emergency vehicles, access roads, shelters, water sources, and so on. Maps can be used to support prevention, preparedness and response to an epidemic.

How to make a community map

If possible, obtain or create a digital map of the community. If not available, it is possible to draw a simple spatial map that shows the community and all its key reference points. While keeping the fundamental principles of data protection, a community map should include the following:


- The whole community: concentrations of people, their houses, and who lives where
- The main shared/public locations in the community, such as schools, health centres, places of worship, water sources, markets, playgrounds and community meeting centres, communal livestock farming and livelihood areas such as cattle kraal, live bird markets, abattoirs, etc.
- The location of people who are most at risk [if you can identify them]
- Where the epidemic started and how it is spreading [if known and possible to identify]
- Health hazards and risks (e.g. improper rubbish disposal sites, large vector breeding sites)

Using the community map

The map can be used to mark new cases and/or referred cases. Do as follows:

- Form teams to cover certain areas of the map.

- Ensuring the participation of members from the community, each team should find out what it can about its area (how many people are sick, who is vulnerable, how many have been referred to health authorities, any other relevant information). If a zoonotic disease outbreak is suspected or implicated, find out who among the community members is keeping animals, how many animals are sick or have died, and/or presence of vectors in households, environment or even in surrounding water bodies. Work with your manager to target and prioritize those who are most at risk. This will require targeting geographically and, within those identified areas, targeting the most in need based on a vulnerability and capacity analysis that includes a gender and diversity analysis.
- Combine the maps of different teams. In doing this, you will be able to see:
 - Which areas of the epidemic you are covering, which areas you may not be covering, and details of each area. This will help you plan your actions. Some of these actions might include: environmental clean-up; distribution of bed nets; immunization campaigns; quarantine, animal biosecurity measures and other activities associated with managing the epidemic.



Making a community map

03. Communicating with the community

Overview

Communicating during an epidemic can be difficult. Disease outbreaks, especially new ones, can cause uncertainty, fear and anxiety that can result in circulation of rumours, disinformation and misinformation. People may not trust the authorities, the health system or organizations including the Red Cross Red Crescent. They may not listen or may not believe the information they receive from people or organizations they do not trust. People may also be overcome with grief for those who are sick or who have died.

Sometimes, communities have strong beliefs that are different from the preventive and protective social measures promoted by the authorities and healthcare providers. They may believe strongly in their own cultural practices, traditional medicine, or other methods that might not prove effective against the disease. They may not accept certain treatments (including medicines and vaccines).

In many countries messages take the form of directives and one-way-communication. However, community engagement and participation have played a critical role in successful disease control and elimination campaigns in many countries

During a disease outbreak, trusted communication with the community is vital. To build trust, two-way communication is important. "Two-way" means volunteers should both *give messages to* AND *receive messages from* the community. Community members must feel respected and listened to and should have the opportunity to share their beliefs, fears and concerns. To accept volunteers' messages, community members must be able to trust you and have confidence in what you say. Once you understand the beliefs, fears and concerns of community members, you can provide them with truthful and accurate messages.

Providing health messages that are consistent, clear and easy to understand also helps to build trust. Giving accurate information to the community is critical, especially when it is necessary to persuade people to adopt safe practices (which might be different from what they would normally do). Some changes in behaviour that may be promoted are things such as:

- Accepting vaccinations or other medical treatments
- Washing hands with soap at crucial times
- Wearing personal protective equipment
- Burying loved ones in ways that are different from what they would normally do (safe and dignified burials)
- Practising social distancing
- Wearing insect repellent or sleeping under bed nets
- Agreeing to be isolated from others to avoid infecting them
- Preparing food and water differently (often by cleaning, boiling or cooking thoroughly)
- Quarantine and culling of animals (which in the case of livestock animals, is often a main source of food, nutrition and livelihood and may be difficult to accept by the farmers who own them)
- And other recommended public health measures

What to do and how to do it

Communicating in an epidemic

- Engage and involve community leaders and community members
 - Find out where the community obtains its information: Who do they trust to give them health

information (for example: health authorities, community or religious leaders, doctors, traditional healers)

- Work with communities to identify, choose and plan appropriate solutions for stopping the spread of disease
- Talk to members of the community about their ideas, fears, beliefs and actions
 - Try to understand how much they know about the disease and its transmission
 - Try to understand beliefs and practices that might impact the spread of the epidemic
 - Try to understand what motivates or helps them to change behaviours
 - Try to understand what stops them from changing their behaviour
- Use different methods of communication
 - Use two-way communication when possible
 - When you understand the community's beliefs, fears and concerns, try to address these in your own messages
 - Sometimes, one-way communication methods are used to spread health messages to large numbers of people quickly
 - One-way communication methods should always be accompanied by two-way communication methods to ensure the community perspectives are known and listened to
 - People learn and retain information differently. It is important to use different methods
 - Communities are composed of different people and groups who may have different communication preferences or needs.
 - Think about how to target different groups, especially those who are hidden, stigmatized or considered "different" because of their religion, sexual orientation, age, disability, illness, or for any other reason:
 - Think about where you will go to reach them
 - Find out if they trust the same or different sources than other groups within the community
 - Discover if they have different access needs, such as language translation or in case of a disability – a different method of communication
 - When choosing methods of communication, consider what people prefer, trust and can access easily
 - Think about the characteristics of your target groups (for example, do they have access to media, such as radio or television? Can they read if they receive pamphlets of information and in what language? Are they accustomed to getting information from social media? Etc.)
 - Think about the resources you have access to (for example: do you have access to poster printing? Is there an appropriate location within the community where you can offer to answer questions or give out information? Etc.)
 - Consider the content of your message(s) and think about the most appropriate way to share that content in the specific context (for example: targeting men and women separately)
- Communication should be:
 - **Simple and short.** People should be able to understand messages easily and be able to remember and repeat them accurately and without difficulty.
 - **Trusted.** Delivered by people the community trusts, by a method the community trusts (for example: radio, television, posters, town-hall discussions, market meetings etc.).
 - **Accurate and specific.** Always provide correct and precise information. Messages should be consistent and should not be cause for confusion. *If* messages must change (due to new and advancing information about the epidemic), be honest and clear about what has changed and why.
 - **Focused on action.** Messages should be action-oriented and should advise members of the community about what they can do to protect themselves and others.
 - **Feasible and realistic.** Make sure that people have the capacity and resources to carry out the actionable advice you give.
 - **Context-specific.** Information should reflect the needs and situation of the specific community. In all your messages, take account of social and cultural factors that might encourage community members to

adopt safer behaviours (such as accepting vaccines) or prevent them from doing so.

Different ways of communicating

There are many, many ways to communicate with communities. The following one and two-way methods of communication are some examples you might consider. Methods can (and should) be combined to ensure accessibility to as many community members as possible.

- One-way communication methods
 - Video, films, television commercials
 - Songs, poems, drama, role-play or theatre, or other edutainment methods
 - Community announcements such as: community town-criers, loud-speaker announcements, SMS or WhatsApp, mass messaging, social media messages, radio broadcasts
 - Posters, billboards
- Two-way communication methods
- Door-to-door visits
- Meeting with key informants such as: community or religious leaders; traditional healers or midwives; teachers; elders, etc.
- Community discussions encouraging participatory methods such as: three pile sorting, voting charts, mapping, polling, barrier analysis, community planning
- Use of feedback and suggestion boxes or presence of trusted focal persons to receive anonymous feedback or messages from community members.

Pay attention to rumours

Rumours can cause panic and fear or can promote unsafe practices. Under the influence of the rumours, communities can lose trust in the health authorities, and they may lose belief in the ability to stop the epidemic. Rumours sometimes cause people to reject interventions that could prevent the spread of disease. Volunteers must:

- Listen for rumours or incorrect information.
 - Note when and where a rumour was heard and report it to your volunteer supervisor or National Society focal point immediately
 - Try to understand why the rumor is spreading fast and of what importance it is to the community. For example, is it just based on lack of knowledge or fear of the unknown, or is it associated with certain socio-cultural beliefs or associated with the stigmatization of a certain demography of people?
- Correct the rumour
 - Give the community clear, simple facts about the disease
 - Reiterate and explain clearly what they can do to protect themselves and others

04. Community referral to health facilities

Overview

During an epidemic, sick people frequently cannot be treated at home or by volunteers or family. They require medical care and need to go for treatment to a health clinic or hospital.

When carrying out epidemic prevention and control activities in the community, always keep the idea of referral in mind.

A community referral is a recommendation (often made by a community volunteer) to seek services at a health facility or from a health care professional. This recommendation is usually based on the identification of signs of disease or the risk that a disease poses to a person, family or community. A community referral is not a confirmation of illness, nor is it a guarantee that any specific treatment will be given. A diagnosis, and any subsequent treatment, is determined by a health professional and not by the community volunteer.

What to do and how to do it

Identifying people who need to be referred

- Learn the symptoms of the disease that is causing the epidemic and the signs that indicate that affected people should be referred to health facilities
- Always keep your own safety and protection in mind
 - With the advice of your supervisor, find out how you can tell when a person is severely ill and needs to be referred

Map and visit referral facilities

1. Unless there is only one referral facility in the community, the selection of a health facility for volunteers to which to send community referrals should be done by a health professional supporting or working at the National Society and validated by the National Society leadership. Volunteers cannot decide alone to which facilities they can send referrals.
2. Once a health facility has been identified and validated by the National Society, with the support of your manager, visit health facilities and talk to doctors and nurses to coordinate referrals
 - Inform them about Red Cross Red Crescent branch activities in which you are involved and how this may lead to community referrals from branch volunteers trained in epidemic prevention and control
 - Discuss the best method for sending sick people from the community to the health facilities:
 - Public transport?
 - Can people access it? Pay for it?
 - Can sick people use it?
 - Is there a risk of disease transmission to other passengers?
 - Ambulance services?
 - Does the health facility have ambulances?
 - Does the Red Cross Red Crescent branch have ambulances?
 - Can people access them? Pay for them?
 - How do you contact the ambulance?
 - Is the disease highly infectious and requiring special transport?

- If the disease is highly infectious (like Ebola or Marburg), special transport must be arranged so that there is no risk that other people could not be infected
- Tell them about your activities and how you plan to do referrals. Take advice from them

Plan and prepare to make referrals

1. Plan how referrals will be made and facilitated
 - Can the National Society provide transport?
 - Do people have money to pay for transport?
 - Does the health facility require prior notification of the referral? If so, how will the health facility be informed of the referral?
2. Always carry the relevant disease tool with you when you are doing community-based referrals
 - This will help you remember what you should know about the disease and its symptoms.

Making a referral

1. Volunteers act on behalf of their National Society and must have the consent of the National Society before undertaking activities. They should be trained in the principles of the Red Cross Red Crescent Movement and should have appropriate training and supervision before making community referrals.
2. Volunteers should obtain the consent of the person to the referral, or of the guardian if it is a child.
3. Volunteers should work to uphold these principles:
 - Confidentiality – It is important to keep information about community members private and not to discuss people's health, healthcare or other private details with others in the community. Remember that breaches of confidentiality often happen unintentionally, for example, when discussing the day's work with friends or family members.
 - Respect – It is important to respect people's choices and decisions, even if you do not agree with their choices.
 - Safety – If you have concerns about the safety or security of a person (in relation to the community referral, or any other aspect of their situation), you should discuss it with your supervisor to find a safe solution if possible.
4. When you refer, always explain clearly to the family concerned what the disease may be, what its symptoms are, and why you think referral is necessary.
 - Give the family information about the health facilities available and how to reach them by different means of transport
 - Help the family if special transportation is needed



Community messages



24. Finding sick people

05. Volunteer protection and safety

Overview

Volunteers work in vulnerable situations and with people of many capacities. Working in epidemics can be risky because volunteers can also catch a disease and fall sick. In addition to physical risks, there may be risks to volunteers' emotional and mental well-being, due to the nature of the work they undertake. It is important to protect from and minimize the impacts of these risks.

Your National Society should provide proper protection for you and other volunteers who are working in epidemics. Your manager is a valuable resource for information and equipment to protect and preserve your physical, emotional and psychosocial wellbeing.

It is important to follow the guidance from your supervisor and National Society and use the level of protection that is appropriate for the situation you are in.

What to do and how to do it

Protecting yourself and others from disease

1. You must be familiar with and trained to use protective equipment before you wear it in an actual disease environment. Try the equipment out beforehand and learn how to use it properly.
 - In certain epidemics like Ebola, Marburg, Lassa fever and plague, full protection should be used whenever you undertake high risk activities. Full protection requires use of personal protective equipment (PPE). (See Action tool [Personal protection equipment \(PPE\) for highly infectious diseases](#))
 - In other epidemics, you should at least use masks and latex gloves and wash hands with soap after contact with an affected person or animal. (See Action tool [Handwashing with soap](#) for instructions in good hand hygiene.)
2. Volunteers should be vaccinated according to country-specific vaccination guidelines (see Action tool [Routine vaccinations](#)).
 - Volunteers should be vaccinated according to the routine vaccination schedule in the country
 - Volunteers may be eligible for vaccination during mass vaccination if applicable
3. Volunteers should be alert to their own physical and psychosocial well-being during an epidemic
 - Volunteers should be alert to stressors in their personal and working lives, and should have a plan in place for how to cope with stress and trauma in a healthy and helpful way
 - This may include stress management techniques that you already use such as exercise, meditation, taking part in hobbies, etc.
 - Your manager is a useful resource for information and tools to use to help you achieve and maintain psychosocial well-being

Understanding common prevention and control measures


Volunteers should learn additional prevention measures for use in epidemics (and before them). These include:

- Vector control measures (see Action tool [Vector control](#))
- Safe handling of animals (Action tool [Handling and slaughtering animals](#))
- Chemoprophylaxis (Action tool [Chemoprophylaxis](#))
- Safe food and water (Action tools [Good food hygiene](#) and [Clean, safe household water](#))

- Hand hygiene (Action tools [Handwashing with soap](#) and [Handwashing in a highly infectious epidemic](#))

Protecting volunteers from harm and liability to others

1. **Volunteers should be protected** if they suffer damage or injury in the course of their work. Accidents can happen, and volunteers can be injured or even killed. Equally, volunteers can harm other people and their property, especially if they have not been properly trained or given the correct equipment.
 - National Societies therefore need to have appropriate insurance policies. Insurance may be needed to pay compensation to volunteers or their families if they are injured or killed; to pay compensation to others if they suffer harm as a result of volunteer actions; and to cover legal costs. The nature of the cover will depend on the legal system in your country. The Movement urges the National Societies to acknowledge and uphold their duty of care towards volunteers, especially if something should happen to them while carrying out their duties. Ask your manager about the type of insurance or coverage through a “safety net” you can receive.
 - Prior to asking volunteers to perform activities that are high-risk (e.g. safe and dignified burials), National Societies should also ensure volunteers have received the necessary vaccinations and protective equipment. What this includes will depend on the context in which you are working and the health policies for staff and volunteers of your National Society.
2. **Volunteers should be informed** of and understand the National Society’s security policy and follow the rules and regulations it sets out. You should also be informed of any changes in the policy and asked to report any incidents of concern.
 - Safety in the community depends on the personal attributes of volunteers, trainers and other team members – how they work together and how they collaborate with people in the community. Volunteers should be culturally sensitive. Your personal behaviour should never cause offence. You should show integrity and should never become a problem for the community. Correct, polite, impartial behaviour is always expected.
 - Volunteers should be proactive in managing and maintaining their own safety and security. This means you should not hesitate to ask your manager about safety and security risks and what you should do if you encounter any threats or have any problems. You should find out what protocols are in place if a safety or security incident occurs, including how and to whom you should report these events.



19. Mental Health and Psychosocial support (MHPSS)

Overview

Normal reactions to abnormal events

It is normal and expected to have strong reactions to abnormal and difficult events. People and communities who experience difficulties may be affected emotionally, mentally, physically and/or socially. Some of these effects may include:

Normal reactions to abnormal events

- **Emotional.** Anxiety, grief, guilt, anger, irritability, frustration, sadness, shame, numbness, loss of hope, loss of meaning, feeling of emptiness.
- **Mental.** Loss of concentration, memory loss, confusion, intrusive thoughts, difficulties in decision making, disorganized thought.
- **Physical.** Increased heartrate, sleeping problems, aches (stomach, head), back and neck pain, muscle tremors and tension, loss of energy, inability to rest and relax.
- **Social.** Risk taking, over- or under-eating, increased intake of alcohol or cigarettes, aggression, withdrawal, isolation.

Psychosocial support

- The term “psychosocial” refers to the dynamic relationship between the psychological and social dimensions of a person, where the dimensions influence each other. The psychological dimension includes emotional and thought processes, feelings and reactions. The social dimension includes relationships, family, community networks, social values and cultural practices.
- “Psychosocial support” refers to actions that meet the psychological and social needs of individuals, families and communities. Psychosocial support (PSS) requires training and supervision. Your supervisor can help you access the appropriate training before you begin to offer PSS to community members. They will also provide you with supervision and support while you provide PSS.
- We provide psychosocial support to help people who have been affected by a crisis. Volunteers should explain what psychosocial support is and if they are appropriately trained, they should offer to provide it to those who wish to receive it. Early and adequate psychosocial support can prevent distress and suffering from turning into more severe mental health problems.
- Psychosocial support during emergencies should ensure safety and promote calm, connectedness, personal and collective efficacy, and hope.

What to do and how to do it

Psychosocial support activities include:

- Psycho-education
 - Explain how to identify signs of psychosocial distress

- Provide advice on how to cope during outbreaks (e.g. maintaining a daily routine as much as possible; calling friends and family to keep in touch and show care for each other; fact-checking information about a disease against trustworthy sources)
- Share tips about relaxation
- Health education can have a positive psychosocial impact:
 - Health education can help community members to better understand their health status, regain a sense of control and cope with their situation
 - While being ill, and even after medical clearance, it can be difficult for people suspected of infection to resume normal life. Educating communities about the nature of the disease, how it spreads – and does not spread – and how to protect against it is an important tool against fear and stigma
- Active listening: Ensure the affected population can raise their concerns, provide suggestions and feedback. This information is used to reduce fear, address rumours and misinformation and increase sense of agency and dignity of the affected population.
- Life skills and vocational skills activities/lessons.
- Creative activities, sports and physical activities.
- Restoring family links.
- Child friendly spaces.
- Supporting memorials and traditional burials.
- Support and self-help groups
 - These include efforts to help people in isolation or quarantine maintain contact with their relatives and friends.
 - Community volunteers that respond to crises are also exposed to loss, devastation, injury and death. It is therefore important to seek support from managers when needed, and to create a supportive environment by showing concern for staff and other volunteers.
- Psychological first aid

36. Vector and reservoir control

Overview


- Vectors are insects or animals that spread infectious diseases through a bite, or contact with their urine, faeces, blood, etc. Many diseases are spread by vectors. Some of these diseases include malaria, dengue fever, Zika, chikungunya, yellow fever, Lassa fever, Rift Valley fever and plague.
- Vectors sometimes live and thrive on other host animals, called reservoirs, before they reach the human population. To protect people from disease, it is important to control both vectors and reservoirs. Vectors and reservoirs include animals and insects such as mosquitoes, ticks, rodents, fleas, etc.

What to do and how to do it

Vector and reservoir control in the community

- Promote hygiene, sanitation and protective practices (** the specific practices you will encourage are dependent on the type of vector or reservoir of concern **). Talk to people in the community about environmental protection strategies against mosquitoes:
 - Repair and close any holes in windows, walls, roofs
 - Use insecticide-treated screens on windows and doors, if available
 - Drain stagnant and standing water, cover water containers
 - Find professionals to spray or larvicide against vectors - spraying chemicals to get rid of vectors can be dangerous, especially if you do not have the proper equipment or materials and do not know how to spray safely. (Only help if you are trained or guided by a trained person.)
 - Outdoor spraying
 - Indoor-residual spraying
- Talk to people in the community about environmental protection strategies against rodents, other small animals and the ticks or fleas that live on them:
 - Store food and water properly, in rodent-proof containers
 - Keep shelters and houses clean
 - Repair and close any holes in windows, walls, roofs
 - Clean the environment of rubbish and waste
 - Keep livestock outside the household (to prevent humans and animals sharing living space)
 - In communities and households infested with rodents, engage environmental health professionals to conduct deratisation exercises
- Talk to people in the community about personal protection strategies against mosquitoes:
 - Use insecticide-treated bed nets to prevent diseases like malaria (not for general use in diseases transmitted by *Aedes* mosquitoes)
 - Wear protective clothing (for example, with long sleeves)
 - Get vaccinated
 - Chemoprophylaxis (preventive treatment)
- Talk to people in the community about personal protection strategies against rodents, other small animals and the ticks or fleas that live on them:
 - Wear protective clothing (for example, with long sleeves)
 - Chemoprophylaxis (preventive treatment)

- Sleep on raised platforms or beds



Community messages



04. Storing water properly



06. Using a clean latrine



07. Protecting yourself against mosquitoes



11. Cleaning up places where mosquitoes breed



12. Good food hygiene



17. Sleeping under mosquito nets



20. Collecting and disposing of rubbish



27. Keeping rodents out

37. Mosquito nets / bed nets

Overview

- Bed nets help to prevent disease spread by mosquitoes, such as malaria. The most effective nets are insecticide-treated nets (ITNs) which are treated with long-lasting insecticides that kills mosquitoes and other insects. The insecticides on these nets are effective for three to five years.
- To be fully effective, nets must be installed and used properly.

What to do and how to do it

Facilitate the use of ITNs


1. Make sure that as many people as possible have bed nets in your community.
- Find out where to obtain ITNs and help your local National Society branch to distribute them.
 - If the Red Cross or Red Crescent is able to distribute nets:
 - Provide a string or piece of rope for people to be able to hang the nets.
 - It is recommended that in each household there is at least one ITN for every two people.

Promote the correct and consistent use of ITNs

1. Encourage every member of the community to sleep under a mosquito net at night and during daytime naps (depending on the type of mosquito and the time of day it typically bites).
 - It is particularly important for children and pregnant women to do so.
2. Show people how to hang their nets properly: Visit households or organize community sessions to demonstrate how to hang nets and encourage their use.
 - Hang the net above the sleeping space.
 - Make sure the net is tucked under the mattress or mat to prevent mosquitoes from getting inside.
 - Keep the net closed while sleeping and during the day while away from the sleeping place.
3. Explain the use of new nets and their care
 - New nets should be hung outside and aired for one day.
 - Care for the net
 - Repair any rips or tears in the net by sewing them closed or using patches
 - Washing the net can remove insecticide. When you wash the net, follow the washing instructions provided with the net. Do not wash the net when it is new and do not wash it too often.
 - Replace the nets at regular intervals, as advised (every three to five years).
4. Follow up to ensure that households use their bed nets.
 - If people are not using their bed nets, find out why.
 - If you see Red Cross or Red Crescent-distributed nets being sold in a local market or used for other purposes (e.g. as construction material or fishing nets), inform your manager.



Community messages



17. Sleeping under mosquito nets

38. Waste disposal and clean-up campaigns

Overview

- Diseases can spread easily in environments where there is waste and lack of sanitation. Rubbish can attract rodents and create breeding and feeding sites for various domestic, wild and scavenging animals and insects (including mosquitoes and flies). Human or animal faeces can also attract flies and other insects that carry germs, cause diseases and contaminate water sources.
- Human and animal waste infected with certain diseases (including animal carcasses, hospital waste, etc.) can also attract different animals and insects which may then spread the diseases to other animals and to humans.
- Standing and stagnant water provide breeding sites for various insects, including mosquitoes.

What to do and how to do it

Planning and preparing for clean-up

- Work with traditional and community leaders, the village health committee and other community partners to decide what needs to be cleaned up and how to go about it.
 - Help to organize activities on agreed “clean-up” days.
 - Ask community leaders to organize volunteer groups for each clean-up initiative.
- Speak with community members about the importance of keeping their community clean.
- Organize a community initiative to plan and create a central garbage disposal area.
- Make sure to have the cleaning tools and supplies available in advance to distribute among the community.

Facilitating community clean-up

- Organize special clean-up days in which all members of the community participate (twice a year or more often if possible).
- Motivate and support the community to work together to:
 - Keep the community free of animal faeces and urine.
 - Keep the community free of puddles and other mosquito breeding sites.
 - Keep the community free of rubbish and garbage (by burning or burying it).
 - Clear vegetation from around riverbanks and ponds near the community
 - Clean the areas around water sources (such as pumps and wells).
 - Create and maintain water soak pits around water sources.



Tell community members how important it is to keep their community clean. Organize regular clean-up campaigns. Include houses and latrines, etc.

What you can do

- Work with traditional and political leaders, the village health committee and other community partners to decide what needs to be cleaned up and how to go about it.
- Help to organize activities on agreed “clean-up” days.
- Ask community leaders to organize volunteer groups for each clean-up initiative.
- Organize special clean-up days in which all members of the community participate (twice a year or more often if possible).
- Organize a community initiative to plan and create a central garbage disposal area.

Motivate and support the community to work together to:

- Keep the community free of animal faeces.
- Keep the community free of puddles and other mosquito breeding sites.
- Keep the community free of rubbish and garbage (by burning or burying it).
- Clean the areas around water sources (such as pumps and wells).
- Create and maintain water soak pits around water sources.



Make sure to dispose of waste properly. This will help to protect the community from germs.

Community messages



07. Protecting yourself against mosquitoes



17. Sleeping under mosquito nets



20. Collecting and disposing of rubbish

42. Promoting safe sex

Overview

Some epidemic-causing diseases can be passed from one person to another during sex, usually in semen, vaginal fluids or blood. Some of these diseases can continue to be spread through sex, even after the person has recovered from the disease. “Sex” means any kind of oral, vaginal or anal sex, or sharing of sex toys. Some epidemic-causing diseases that are known to be passed through sex are:

- Zika virus
- Ebola virus disease
- Marburg fever
- Lassa fever
- Mpox virus

There are many other diseases, such as HIV/AIDS, which can be passed through sexual activities. Promoting safer sex is an important public health message that can save many lives. Having “safer sex” means protecting yourself and your partner from diseases that can be transmitted during sexual activity.

When to promote safer sex?

During outbreaks of Zika, Ebola, Marburg fever, mpox or Lassa fever, it is important to include messaging about safer sex. Messages should include:

- How the disease is transmitted through sex
- How to practise safer sex

While safer sex messaging is very important to prevent the spread of epidemic disease, it is only one way that community members should protect themselves from becoming ill during outbreaks of the above diseases. Especially in the cases of Ebola, Marburg, mpox and Lassa fevers, the diseases are also spread through other close contact with infected people, not just sex. Practising safer sex alone will not protect people from these diseases. Safer sex is only one way to prevent disease. It should not be the principal focus of your messages during an epidemic.

Messages about safer sex during epidemics should promote:

- Use of male or female condoms when having sex.
 - Condoms are a barrier and block any infectious semen or fluid that may transmit an infection.
- Other ways (instead of sexual intercourse) of sharing intimacy with sexual partners

Key facts about Zika:

- A pregnant woman can pass Zika to her unborn baby which can cause severe brain defects in the baby
 - Pregnant women who live in or travel to places where Zika is present, or whose sexual partners live in or travel to places where Zika is present, should be instructed to go to a health centre for a check-up and to discuss Zika risk with their healthcare provider.
 - In these cases, it is best not to have sex during pregnancy, or to use condoms during the whole pregnancy.
- If women are planning a pregnancy and live in an area where Zika is present, it is important to discuss the risks, protect both partners from mosquito bites, and consider postponing pregnancy until after the

outbreak has ended

- A man or woman who has Zika can pass Zika to his or her partner during sex for up to six months after acquiring the infection (whether they were sick and showed symptoms or not).

Key facts about Ebola, Marburg and Lassa:

- Men who have recovered from Ebola, Marburg fever or Lassa fever can pass the disease on to another person during oral, vaginal or anal sex
- Men who have recovered from Ebola or Marburg should use a condom for at least 12 months from when they got sick or until their semen tests negative twice for the virus.
- Men who have recovered from Lassa fever should use a condom for at least three months after they get better.

Key facts about mpox

- During the global outbreak that began in 2022, the virus mostly spread through sexual contact. Anyone who has close contact with someone who has symptoms and any person with multiple sex partners is also at risk. Therefore, people who have multiple or new sexual partners including those who are sexually active or gay, bisexual and other men who have sex with men may be at higher risk of being exposed if they have sex or other form of close contact with someone who is infectious.
- Therefore, it is important to engage communities including those of sex workers, gay, bisexual and other men who have sex with men to raise awareness on risks of mpox.


What to do and how to do it

Social mobilization, messaging, and behaviour change

1. Make sure you understand the facts, and how diseases including Zika, Ebola, Marburg fever, mpox and Lassa fever can be transmitted through sex.
2. Remember that sex is only one way of transmitting these diseases, and often it is not the main way.
 - Talk to your volunteer coordination/team leader or local branch health office about which prevention methods should be emphasized during your community visits.
3. Remember that sex can be a very sensitive, taboo or embarrassing topic for people to talk about. You may have to change your approach or way of communicating to get your messages across.
4. Carry out social mobilization and behaviour change communication activities in an outbreak of Zika, Ebola, Marburg fever, mpox or Lassa fever. (See Important points above and Action Tools [*Communicating with the community*](#) and [*Social mobilization and behaviour change*](#).)
 - Make sure you know and can demonstrate how to use a male and female condom correctly.
5. Be respectful of the culture:
 - Consider speaking to men and women separately
 - Consider speaking separately to adolescents. Remember that adolescents often experience stigma and difficulties in accessing sexual health information and services. Yet, most people initiate sexual activity during adolescence, so it is important to support them to ensure good sexual health choices and decisions.
 - Consider adapting specific communication strategies and schedules to key populations as they are often marginalized and even hard-to-reach due to stigmatization or unfavorable laws in some countries.
 - Do not push people to share views on this topic in front of others as some may feel uncomfortable
 - Do not make assumptions based on stereotypes. For example, do not assume a person has certain

attitudes about sex based on their age, gender, profession, or ethnicity

- Remember that providing sexual health information is important, even if it is a sensitive topic in many cultures



Community messages



26. Practising safe sex

43. Social mobilization and behaviour change

Overview

There are many reasons why people practise unhealthy behaviours. People are affected by access to services or facilities, social norms and influences where they work, live or play. Behaviour change is the study of how and why people change some habit or action in their life. As volunteers, we need to understand WHY the behaviour is happening and WHAT actions will lead to change to create healthy behaviours. Examples of healthy behaviours include handwashing, breastfeeding, immunizations, consistent condom use and use of bed nets.


In any culture and context, behaviour change involves three elements. Before people will change their behaviour:

1. They need to know what, why and how they should change. They need knowledge.
2. They need to have the right equipment, access and capacity. They need an enabling environment.
3. They need to be motivated to change.



The social-ecological model below shows how each person's behaviours are affected by many different levels of influence including the individual level, the interpersonal level, the community level, the organizational level and the broader policy level which includes laws and policies that allow or restrict a behaviour. In order to promote health, it is important to consider and plan behaviour change activities across multiple levels at the

same time. This approach is more likely to result in successful behaviour change over time. As a volunteer, it is helpful to understand that behaviour change is difficult for many people because of these many levels and the complex interactions and expectations across the different levels. If you consider how each of the levels affects the behaviour of the person you want to help, you can try different interventions at each level that is specific to their needs.



What do to and how to do it

The general process for developing a behaviour change intervention includes staff and volunteers working through the general steps of:

1. Sensitizing the community to the behaviour change process using the theory of change model.
2. Assessing the problem behaviour – why it is practised, who practises it, when it is practised and what factors in the environment or society encourage the behaviour. Assess this information at the different levels of the social-ecological model for each community you serve.
3. Identifying an appropriate behaviour goal based on your assessment.
4. Reviewing the causes or barriers at each level that allow the behaviour to continue. Identify interventions that align with each cause or barrier and that can be used at different levels.
5. Discussing the suggested interventions for each social-ecological model level with the community.
6. Identifying appropriate interventions for the context at each level. Interventions should be planned to address the stages of the theory of change by first giving knowledge and addressing environmental factors, motivating key people to gain approval and intentions, and ultimately inciting people to action that contributes to the overall goal.
7. Implementing the interventions at each level.

8. Monitoring to see if change is happening. Change takes time but it must be monitored to ensure that it is happening, even slowly. Additionally, as people go through the change process, their barriers and causes will change. The behaviour change interventions should adjust to these changes to ensure that change can continue.
9. Recognizing that when change is not happening as intended, further assessment and intervention tweaking is needed.
10. Continuing to implement, monitor, assess and adjust as the change process happens.

For more information, please consult the eCBHFA Manual for volunteers on [Behaviour Change](#), including:

1. Principles of behaviour change
2. The social ecological model
3. The stages of behaviour change
4. Activities for behaviour change

Community messages



23. Encouraging healthy behaviours in a community