



# Ebola virus disease (EVD)

Last update: 2025-03-18

## Key facts

- Ebola virus disease (EVD or Ebola) is a rare but severe and often fatal illness in humans.
- The average Ebola case fatality rate is around 50%, and this has varied from 25–90% in past outbreaks, depending on circumstances and the response.
- Ebola first appeared in 1976 in 2 simultaneous outbreaks in South Sudan, and the Democratic Republic of Congo. The latter occurred in a village near the Ebola River, from which the disease takes its name.
- The origin of the virus is unknown, but current evidence suggests that fruit bats (Pteropodidae) may be a host.

### Transmission: Direct contact

Ebola is introduced into the human population through close contact with the blood, secretions, organs or other bodily fluids of infected animals such as fruit bats, chimpanzees, gorillas, monkeys, forest antelope or porcupines found ill or dead or in the rainforest. This may happen by:

- Touching or eating animals that are infected (monkeys, apes, bats)
- This is especially true if bushmeat is not well cooked
- Touching the waste (faeces/poop) from infected fruit bats
- Through direct contact with tissue, blood, other body fluids, and excretions from an infected animal.

Ebola then spreads through human-to-human transmission through the following ways:

- Contact with the blood, faeces/poop, vomit, spit, sweat, tears, breast milk or semen from a person sick with Ebola
  - Contact can be with objects that have been contaminated with the body fluids from a person who is sick with Ebola (for example, bedding, clothes or needles)
- When handling the body of someone who has died from Ebola
- Coming into contact with semen (for example, through sex) from someone who has recovered from Ebola (Ebola can be spread through semen for up to 12 months after recovery)
- Pregnant women who get acute Ebola and recover from the disease may still carry the virus in breastmilk, or in pregnancy related fluids and tissues.

## Most vulnerable to severe consequences

- Every person in the community is vulnerable to Ebola (\*\* it is a very deadly disease than can kill many – half to almost all people who get infected can die \*\*)
- Pregnant and breastfeeding women

## Most vulnerable to contracting the disease

- Family and relatives of people who are sick (or who have died) from Ebola
- Health workers, volunteers, and people who handle dead bodies

## Symptoms

At the beginning there may be:

- Sudden fever
- Tiredness
- Muscle pain
- Headache
- Sore throat

Followed by:

- Vomiting
- Diarrhoea
- Rash

Bleeding - under the skin or from the nose and gums (inside mouth), coughing blood or having bloody faeces (sometimes)

**Marburg and Ebola viruses which are both members of the same virus family, have similar clinical signs, and have the capacity to cause outbreaks with high fatality rates**

# What can you do to prevent and control an epidemic?

## Monitoring the community and identifying sick people

- Identify and ensure that sick people immediately isolate

## Treatment and management

- Immediately refer suspected cases to treatment centres
  - Provide safe transport for suspected cases
- Support contact tracing and follow-up of contacts
- Provide psychosocial support to the sick person, their families, and communities
- Conduct safe and dignified funeral and burial practices

## Sanitation and waste management

- Dispose safely of waste that might be contaminated (by burning or burying)
- Disinfect homes and personal belongings of people who are sick or have died
  - Avoid disinfection in spaces where those who are sick or have died have not been present. Disinfection in such places can create stigma unnecessarily, is not an effective prevention method, and is an inappropriate use of resources.
- Promote disinfection of reusable supplies
- Solid waste management
  - Encourage the community to clear away rubbish and garbage

## Personal protection and hygiene

- Promote handwashing with soap, chlorine solution or hand sanitizer
- Practise and promote social distancing
- Use personal protective equipment (gloves, masks, clothing) when in contact with potential cases
- Avoid touching sick or dead fruit bats, monkeys, apes and other bush meat
- Avoid contact with the faeces and other waste products of bushmeat

## Food hygiene and safety

- Cooking bushmeat or animal products (blood and meat) thoroughly

## Social mobilization and health promotion

- Find out the specific advice being given by health and other relevant authorities

- Promote the use of condoms for at least 12 months after a diagnosis of Ebola
- Promote safe and dignified burials
- Promote the safe disposal (burning or burying) of contaminated materials
- Promote social distancing
- Model following this advice and inform community members of current health practice advice
- Offer support and encouragement to follow the advice
  - Try to gain understanding about if and why health practice advice is not being followed
  - With the guidance of your supervisor and health authorities, work with communities to overcome barriers to following health advice and recommended practices
- Involve the community in managing rumours and misinformation

## Immunization

- Support vaccination campaigns where available

# Mapping and community assessment

- Make a map of the community.
- Mark the following information on the map:
  - How many people have fallen sick with Ebola? Where?
  - How many people have died? Where? When?
  - Where are the local health facilities and services? (include traditional healers)
- Record the following information on the back of the map:
  - When did people start to fall sick with Ebola?
  - How many people live in the affected community?
  - What are the community's habits, practices and beliefs about caring for sick people? Consider any differences in roles and responsibilities between women and men.
  - What are the community's burial traditions, funeral procedures and practices? Who traditionally prepares bodies for burials? Consider any differences in roles and responsibilities between women and men.
  - Do people in the community know about Ebola?
    - Do people know the main signs of Ebola?
    - Do they know what to do if someone becomes sick (for example phone number to call, actions to take)?
    - Do people know how to protect themselves from Ebola?
  - Is a social mobilization or health promotion programme in place?
  - Are people in the community socially distancing? Why? Why not?
  - Which sources do people use/trust the most for information?
    - Are there rumours or misinformation about Ebola? What are the rumours?
  - Are health workers, volunteers or people who have survived Ebola stigmatized, left out, threatened or harassed?

- What are the main effects on them and their lives?

## Volunteer actions

- [01. Community-based surveillance](#)
- [02. Community mapping](#)
- [03. Communicating with the community](#)
- [04. Community referral to health facilities](#)
- [05. Volunteer protection and safety](#)
- [06. Personal protection equipment \(PPE\) for highly infectious diseases](#)
- [19. Mental Health and Psychosocial support \(MHPSS\)](#)
- [20. Isolating sick people](#)
- [25. Mass vaccination campaigns](#)
- [28. Physical distancing](#)
- [34. Handwashing with soap](#)
- [35. Handwashing in a highly infectious epidemic](#)
- [41. Handling and slaughtering animals](#)
- [42. Promoting safe sex](#)
- [43. Social mobilization and behaviour change](#)

## Other resources

### Additional References

[Ebola Virus Disease - WOA - World Organisation for Animal Health](#)

[WHO – World Health Organization; Ebola Virus Disease; 2023](#)

[WHO – World Health Organization; Ebola Virus Disease, Question and Answer; 2019](#)

[WOAH – World Organization for Animal Health; Ebola Virus Disease, 2024](#)

# 01. Community-based surveillance

## Overview

- Community-based surveillance is the systematic detection and reporting of significant public health events (such as sudden illness or death in people or animals) within a community by community members and volunteers. It is a simple, adaptable, low-cost public health initiative designed to complement early warning systems for potential epidemic diseases.
- Volunteers use something called a “community-case definition” to detect and report signs and symptoms of potential diseases, health risks and events, and support in community actions and response of local health authorities. Community case definitions are designed to align with the local language and do not require medical training to report on.
- Information discovered during surveillance should be shared with the local branch and health authorities based on the agreed protocol. Where relevant, (e.g. for zoonoses or environmental health events) information should also be shared with animal health and environmental health authorities.
- Community-based surveillance (CBS) can be done alongside other health, WASH or community engagement activities in your community, and therefore is not a stand-alone activity, but one that is useful to partner with other community-based activities.
- Community-based surveillance aids in
  - Early detection of public health risks within the community
  - Complementing early warning systems, extending them to the community
  - Linking early detection to early action within the community

## What to do and how to do it

- Preparation activities
  - Work with supervisors in mapping community needs and human, animal and environmental disease priorities (see Action tool [Community mapping](#))
  - Familiarize yourself with the disease that may be present in your community including signs and symptoms
  - Establish who is vulnerable in the community. Doing this will help you to identify people who are more likely to fall sick
  - Ensure referral mechanisms are clear in case community members fall sick and require referrals to health facilities for care.
  - Engage in community engagement activities such as mobile cinema, house-to-house visits, etc. to remain active and a known resource in the community.
- Recognize
  - Detect signs and symptoms corresponding to human, animal or environmental health risks or events in your community aligned with community case definitions
  - When you detect people who are sick with the disease, assess how severely ill they are and whether they need to be referred to a health facility (see Action tool [Referral to health facilities](#)).
  - Record the health risk or event you detected to ensure it can be followed-up

- Report
  - Report on the detected health risks or event in your community to your supervisor based on the methodology you are trained on (for example, SMS, phone call, or mobile application). *Remember that reporting must be systematic. To avoid confusion, everyone who reports should follow the same methods agreed on in the protocol and in the training.*
  - Your supervisor will then cross-check the report ensuring it meets the community case definition or unusual event requirements agreed on with health authorities. If matching, the supervisor will escalate the alert to the local health authorities for a response or investigation
  - After verification, the supervisor will notify relevant authorities in animal and environmental health for significant animal, zoonotic and environmental health events, especially those that portend a risk to human health.
- React
  - Begin community-level activities based on the health risk following proper safety precautions
  - Referral or care at home
    - Communicate specific health messages and information, and refer sick people promptly to health facilities
    - If sick people can be cared for at home, show their families what to do and provide them with information and supplies, where possible. Use corresponding “volunteer actions” in the ECV toolkit corresponding to the suspected epidemic risk.
  - Support health authorities in their investigation or response following-up on the alert
  - Where relevant, collaborate with and support officials in the animal and environmental health sectors for joint investigation, response and information sharing.

Additional resources on community-based surveillance: <https://cbs.ifrc.org/>

## Community messages



24. Finding sick people

## 02. Community mapping

### Overview

A map of the community enables you to connect issues or problems with particular places and makes information easy to see. Maps are often easier to understand than words.

Mapping aids in:

- Identifying risks and exposure to risk
  - Who and what are most exposed
  - What are they exposed to
- Show existing problems and vulnerabilities (some might make the current threat more serious)
- Understanding resources within the community that might be useful in managing the epidemic
- Obtaining information about other sectors (such as livelihoods, shelter, WASH, infrastructure etc.) that might be influenced by the epidemic, or that might be useful in managing the epidemic
- Analysing links and patterns in the exposure and spread of the epidemic - which may include human-to-human transmission dynamics, exposure from animals, vectors or food, behavioural risks, and environmental health drivers.

It is important to create the map together with community members. This helps communities to be active and to be participating members in the care offered by the Red Cross Red Crescent and volunteers.

Community mapping is especially useful in epidemics because it helps to see where the biggest problems and needs are and helps to identify risks and resources such as health posts, emergency vehicles, access roads, shelters, water sources, and so on. Maps can be used to support prevention, preparedness and response to an epidemic.

### How to make a community map

If possible, obtain or create a digital map of the community. If not available, it is possible to draw a simple spatial map that shows the community and all its key reference points. While keeping the fundamental principles of data protection, a community map should include the following:


- The whole community: concentrations of people, their houses, and who lives where
- The main shared/public locations in the community, such as schools, health centres, places of worship, water sources, markets, playgrounds and community meeting centres, communal livestock farming and livelihood areas such as cattle kraal, live bird markets, abattoirs, etc.
- The location of people who are most at risk [if you can identify them]
- Where the epidemic started and how it is spreading [if known and possible to identify]
- Health hazards and risks (e.g. improper rubbish disposal sites, large vector breeding sites)



## Using the community map

The map can be used to mark new cases and/or referred cases. Do as follows:

- Form teams to cover certain areas of the map.
  - Ensuring the participation of members from the community, each team should find out what it can about its area (how many people are sick, who is vulnerable, how many have been referred to health authorities, any other relevant information). If a zoonotic disease outbreak is suspected or implicated, find out who among the community members is keeping animals, how many animals are sick or have died, and/or presence of vectors in households, environment or even in surrounding water bodies. Work with your manager to target and prioritize those who are most at risk. This will require targeting geographically and, within those identified areas, targeting the most in need based on a vulnerability and capacity analysis that includes a gender and diversity analysis.
- Combine the maps of different teams. In doing this, you will be able to see:
  - Which areas of the epidemic you are covering, which areas you may not be covering, and details of each area. This will help you plan your actions. Some of these actions might include: environmental clean-up; distribution of bed nets; immunization campaigns; quarantine, animal biosecurity measures and other activities associated with managing the epidemic.



Making a community map

## 03. Communicating with the community

### Overview

Communicating during an epidemic can be difficult. Disease outbreaks, especially new ones, can cause uncertainty, fear and anxiety that can result in circulation of rumours, disinformation and misinformation. People may not trust the authorities, the health system or organizations including the Red Cross Red Crescent. They may not listen or may not believe the information they receive from people or organizations they do not trust. People may also be overcome with grief for those who are sick or who have died.

Sometimes, communities have strong beliefs that are different from the preventive and protective social measures promoted by the authorities and healthcare providers. They may believe strongly in their own cultural practices, traditional medicine, or other methods that might not prove effective against the disease. They may not accept certain treatments (including medicines and vaccines).

In many countries messages take the form of directives and one-way-communication. However, community engagement and participation have played a critical role in successful disease control and elimination campaigns in many countries

During a disease outbreak, trusted communication with the community is vital. To build trust, two-way communication is important. "Two-way" means volunteers should both *give messages to* AND *receive messages from* the community. Community members must feel respected and listened to and should have the opportunity to share their beliefs, fears and concerns. To accept volunteers' messages, community members must be able to trust you and have confidence in what you say. Once you understand the beliefs, fears and concerns of community members, you can provide them with truthful and accurate messages.

Providing health messages that are consistent, clear and easy to understand also helps to build trust. Giving accurate information to the community is critical, especially when it is necessary to persuade people to adopt safe practices (which might be different from what they would normally do). Some changes in behaviour that may be promoted are things such as:

- Accepting vaccinations or other medical treatments
- Washing hands with soap at crucial times
- Wearing personal protective equipment
- Burying loved ones in ways that are different from what they would normally do (safe and dignified burials)
- Practising social distancing
- Wearing insect repellent or sleeping under bed nets
- Agreeing to be isolated from others to avoid infecting them
- Preparing food and water differently (often by cleaning, boiling or cooking thoroughly)
- Quarantine and culling of animals (which in the case of livestock animals, is often a main source of food, nutrition and livelihood and may be difficult to accept by the farmers who own them)
- And other recommended public health measures

## What to do and how to do it

### *Communicating in an epidemic*

- Engage and involve community leaders and community members
  - Find out where the community obtains its information: Who do they trust to give them health information (for example: health authorities, community or religious leaders, doctors, traditional healers)
  - Work with communities to identify, choose and plan appropriate solutions for stopping the spread of disease
  - Talk to members of the community about their ideas, fears, beliefs and actions
    - Try to understand how much they know about the disease and its transmission
    - Try to understand beliefs and practices that might impact the spread of the epidemic
    - Try to understand what motivates or helps them to change behaviours
    - Try to understand what stops them from changing their behaviour
- Use different methods of communication
  - Use two-way communication when possible
    - When you understand the community's beliefs, fears and concerns, try to address these in your own messages
  - Sometimes, one-way communication methods are used to spread health messages to large numbers of people quickly
    - One-way communication methods should always be accompanied by two-way communication methods to ensure the community perspectives are known and listened to
  - People learn and retain information differently. It is important to use different methods
    - Communities are composed of different people and groups who may have different communication preferences or needs.
      - Think about how to target different groups, especially those who are hidden, stigmatized or considered "different" because of their religion, sexual orientation, age, disability, illness, or for any other reason:
        - Think about where you will go to reach them
        - Find out if they trust the same or different sources than other groups within the community
        - Discover if they have different access needs, such as language translation or in case of a disability – a different method of communication
    - When choosing methods of communication, consider what people prefer, trust and can access easily
      - Think about the characteristics of your target groups (for example, do they have access to media, such as radio or television? Can they read if they receive pamphlets of information and in what language? Are they accustomed to getting information from social media? Etc.)
      - Think about the resources you have access to (for example: do you have access to poster printing? Is there an appropriate location within the community where you can offer to answer questions or give out information? Etc.)
      - Consider the content of your message(s) and think about the most appropriate way to share that content in the specific context (for example: targeting men and women separately)
- Communication should be:
  - **Simple and short.** People should be able to understand messages easily and be able to remember and repeat them accurately and without difficulty.
  - **Trusted.** Delivered by people the community trusts, by a method the community trusts (for example:

radio, television, posters, town-hall discussions, market meetings etc.).

- **Accurate and specific.** Always provide correct and precise information. Messages should be consistent and should not be cause for confusion. *If* messages must change (due to new and advancing information about the epidemic), be honest and clear about what has changed and why.
- **Focused on action.** Messages should be action-oriented and should advise members of the community about what they can do to protect themselves and others.
- **Feasible and realistic.** Make sure that people have the capacity and resources to carry out the actionable advice you give.
- **Context-specific.** Information should reflect the needs and situation of the specific community. In all your messages, take account of social and cultural factors that might encourage community members to adopt safer behaviours (such as accepting vaccines) or prevent them from doing so.

### ***Different ways of communicating***

There are many, many ways to communicate with communities. The following one and two-way methods of communication are some examples you might consider. Methods can (and should) be combined to ensure accessibility to as many community members as possible.

- One-way communication methods
  - Video, films, television commercials
  - Songs, poems, drama, role-play or theatre, or other edutainment methods
  - Community announcements such as: community town-criers, loud-speaker announcements, SMS or WhatsApp, mass messaging, social media messages, radio broadcasts
  - Posters, billboards
- Two-way communication methods
- Door-to-door visits
- Meeting with key informants such as: community or religious leaders; traditional healers or midwives; teachers; elders, etc.
- Community discussions encouraging participatory methods such as: three pile sorting, voting charts, mapping, polling, barrier analysis, community planning
- Use of feedback and suggestion boxes or presence of trusted focal persons to receive anonymous feedback or messages from community members.

### ***Pay attention to rumours***

Rumours can cause panic and fear or can promote unsafe practices. Under the influence of the rumours, communities can lose trust in the health authorities, and they may lose belief in the ability to stop the epidemic. Rumours sometimes cause people to reject interventions that could prevent the spread of disease. Volunteers must:

- Listen for rumours or incorrect information.
  - Note when and where a rumour was heard and report it to your volunteer supervisor or National Society focal point immediately
  - Try to understand why the rumor is spreading fast and of what importance it is to the community. For example, is it just based on lack of knowledge or fear of the unknown, or is it associated with certain socio-cultural beliefs or associated with the stigmatization of a certain demography of people?
- Correct the rumour

- Give the community clear, simple facts about the disease
- Reiterate and explain clearly what they can do to protect themselves and others

## 04. Community referral to health facilities

### Overview

During an epidemic, sick people frequently cannot be treated at home or by volunteers or family. They require medical care and need to go for treatment to a health clinic or hospital.

When carrying out epidemic prevention and control activities in the community, always keep the idea of referral in mind.

A community referral is a recommendation (often made by a community volunteer) to seek services at a health facility or from a health care professional. This recommendation is usually based on the identification of signs of disease or the risk that a disease poses to a person, family or community. A community referral is not a confirmation of illness, nor is it a guarantee that any specific treatment will be given. A diagnosis, and any subsequent treatment, is determined by a health professional and not by the community volunteer.

### What to do and how to do it

#### *Identifying people who need to be referred*

- Learn the symptoms of the disease that is causing the epidemic and the signs that indicate that affected people should be referred to health facilities
- Always keep your own safety and protection in mind
  - With the advice of your supervisor, find out how you can tell when a person is severely ill and needs to be referred

#### *Map and visit referral facilities*

1. Unless there is only one referral facility in the community, the selection of a health facility for volunteers to which to send community referrals should be done by a health professional supporting or working at the National Society and validated by the National Society leadership. Volunteers cannot decide alone to which facilities they can send referrals.
2. Once a health facility has been identified and validated by the National Society, with the support of your manager, visit health facilities and talk to doctors and nurses to coordinate referrals
  - Inform them about Red Cross Red Crescent branch activities in which you are involved and how this may lead to community referrals from branch volunteers trained in epidemic prevention and control
  - Discuss the best method for sending sick people from the community to the health facilities:
    - Public transport?
      - Can people access it? Pay for it?
      - Can sick people use it?
      - Is there a risk of disease transmission to other passengers?
    - Ambulance services?
      - Does the health facility have ambulances?

- Does the Red Cross Red Crescent branch have ambulances?
- Can people access them? Pay for them?
- How do you contact the ambulance?
- Is the disease highly infectious and requiring special transport?
  - If the disease is highly infectious (like Ebola or Marburg), special transport must be arranged so that there is no risk that other people could not infected
  - Tell them about your activities and how you plan to do referrals. Take advice from them

### ***Plan and prepare to make referrals***

1. Plan how referrals will be made and facilitated
  - Can the National Society provide transport?
  - Do people have money to pay for transport?
  - Does the health facility require prior notification of the referral? If so, how will the health facility be informed of the referral?
2. Always carry the relevant disease tool with you when you are doing community-based referrals
  - This will help you remember what you should know about the disease and its symptoms.

### ***Making a referral***

1. Volunteers act on behalf of their National Society and must have the consent of the National Society before undertaking activities. They should be trained in the principles of the Red Cross Red Crescent Movement and should have appropriate training and supervision before making community referrals.
2. Volunteers should obtain the consent of the person to the referral, or of the guardian if it is a child.
3. Volunteers should work to uphold these principles:
  - Confidentiality – It is important to keep information about community members private and not to discuss people's health, healthcare or other private details with others in the community. Remember that breaches of confidentiality often happen unintentionally, for example, when discussing the day's work with friends or family members.
  - Respect – It is important to respect peoples' choices and decisions, even if you do not agree with their choices.
  - Safety – If you have concerns about the safety or security of a person (in relation to the community referral, or any other aspect of their situation), you should discuss it with your supervisor to find a safe solution if possible.
4. When you refer, always explain clearly to the family concerned what the disease may be, what its symptoms are, and why you think referral is necessary.
  - Give the family information about the health facilities available and how to reach them by different means of transport
    - Help the family if special transportation is needed





## Community messages



24. Finding sick people



## 05. Volunteer protection and safety

### Overview

Volunteers work in vulnerable situations and with people of many capacities. Working in epidemics can be risky because volunteers can also catch a disease and fall sick. In addition to physical risks, there may be risks to volunteers' emotional and mental well-being, due to the nature of the work they undertake. It is important to protect from and minimize the impacts of these risks.

Your National Society should provide proper protection for you and other volunteers who are working in epidemics. Your manager is a valuable resource for information and equipment to protect and preserve your physical, emotional and psychosocial wellbeing.

It is important to follow the guidance from your supervisor and National Society and use the level of protection that is appropriate for the situation you are in.

### What to do and how to do it

#### *Protecting yourself and others from disease*

1. You must be familiar with and trained to use protective equipment before you wear it in an actual disease environment. Try the equipment out beforehand and learn how to use it properly.
  - In certain epidemics like Ebola, Marburg, Lassa fever and plague, full protection should be used whenever you undertake high risk activities. Full protection requires use of personal protective equipment (PPE). (See Action tool [Personal protection equipment \(PPE\) for highly infectious diseases](#))
  - In other epidemics, you should at least use masks and latex gloves and wash hands with soap after contact with an affected person or animal. (See Action tool [Handwashing with soap](#) for instructions in good hand hygiene.)
2. Volunteers should be vaccinated according to country-specific vaccination guidelines (see Action tool [Routine vaccinations](#)).
  - Volunteers should be vaccinated according to the routine vaccination schedule in the country
  - Volunteers may be eligible for vaccination during mass vaccination if applicable
3. Volunteers should be alert to their own physical and psychosocial well-being during an epidemic
  - Volunteers should be alert to stressors in their personal and working lives, and should have a plan in place for how to cope with stress and trauma in a healthy and helpful way
    - This may include stress management techniques that you already use such as exercise, meditation, taking part in hobbies, etc.
    - Your manager is a useful resource for information and tools to use to help you achieve and maintain psychosocial well-being

#### *Understanding common prevention and control measures*

Volunteers should learn additional prevention measures for use in epidemics (and before them). These include:

- Vector control measures (see Action tool [Vector control](#))
- Safe handling of animals (Action tool [Handling and slaughtering animals](#))
- Chemoprophylaxis (Action tool [Chemoprophylaxis](#))
- Safe food and water (Action tools [Good food hygiene](#) and [Clean, safe household water](#))
- Hand hygiene (Action tools [Handwashing with soap](#) and [Handwashing in a highly infectious epidemic](#))

## Protecting volunteers from harm and liability to others

1. **Volunteers should be protected** if they suffer damage or injury in the course of their work. Accidents can happen, and volunteers can be injured or even killed. Equally, volunteers can harm other people and their property, especially if they have not been properly trained or given the correct equipment.
  - National Societies therefore need to have appropriate insurance policies. Insurance may be needed to pay compensation to volunteers or their families if they are injured or killed; to pay compensation to others if they suffer harm as a result of volunteer actions; and to cover legal costs. The nature of the cover will depend on the legal system in your country. The Movement urges the National Societies to acknowledge and uphold their duty of care towards volunteers, especially if something should happen to them while carrying out their duties. Ask your manager about the type of insurance or coverage through a “safety net” you can receive.
  - Prior to asking volunteers to perform activities that are high-risk (e.g. safe and dignified burials), National Societies should also ensure volunteers have received the necessary vaccinations and protective equipment. What this includes will depend on the context in which you are working and the health policies for staff and volunteers of your National Society.
2. **Volunteers should be informed** of and understand the National Society's security policy and follow the rules and regulations it sets out. You should also be informed of any changes in the policy and asked to report any incidents of concern.
  - Safety in the community depends on the personal attributes of volunteers, trainers and other team members – how they work together and how they collaborate with people in the community. Volunteers should be culturally sensitive. Your personal behaviour should never cause offence. You should show integrity and should never become a problem for the community. Correct, polite, impartial behaviour is always expected.
  - Volunteers should be proactive in managing and maintaining their own safety and security. This means you should not hesitate to ask your manager about safety and security risks and what you should do if you encounter any threats or have any problems. You should find out what protocols are in place if a safety or security incident occurs, including how and to whom you should report these events.



## 06. Personal protection equipment (PPE) for highly infectious diseases

### Overview

When working with certain epidemic diseases, especially highly infectious diseases such as Ebola, Marburg, Lassa fever and plague, it is vital to protect yourself when you touch affected people, their body fluids, or dead bodies or animals.

Like volunteers, PPE comes in different sizes and fits. Not every size of PPE will fit every volunteer appropriately (e.g. protective goggles may not be the right size or shape for some women or for people of certain ethnicities, leaving them at greater risk of exposure). It is very important that volunteers have the correct size and fit of all their equipment for optimum protection.


This tool shows you how to put on and take off PPE. \*\* Note: Different PPE types are used in different contexts, depending on the disease and the type of protection required (e.g. the components of the PPE for Ebola and for plague may have some commonalities but are not the same). Always ensure that you have been properly trained to use the type of PPE available in your context, for the specific disease you are addressing. \*\*

See Action tool [Volunteer protection and safety](#) for more information on volunteer protection.


### What to do and how to do it

- Test the size and fit of your PPE by having your manager check the fit and coverage of all your equipment. If anything is incorrectly sized, be sure you obtain equipment of the correct fit before you begin work that puts you at risk of exposure.

### Steps for putting on protective clothing



### Steps for taking off protective clothing



**Note:** Handwashing, with gloved hands, should be performed before starting to remove protective clothing (step 4), between removing protective clothing but before removing protective face mask (step 9) and at the end with ungloved hands (step 12).

## 19. Mental Health and Psychosocial support (MHPSS)

### Overview

#### *Normal reactions to abnormal events*

It is normal and expected to have strong reactions to abnormal and difficult events. People and communities who experience difficulties may be affected emotionally, mentally, physically and/or socially. Some of these effects may include:

#### Normal reactions to abnormal events

- **Emotional.** Anxiety, grief, guilt, anger, irritability, frustration, sadness, shame, numbness, loss of hope, loss of meaning, feeling of emptiness.
- **Mental.** Loss of concentration, memory loss, confusion, intrusive thoughts, difficulties in decision making, disorganized thought.
- **Physical.** Increased heartrate, sleeping problems, aches (stomach, head), back and neck pain, muscle tremors and tension, loss of energy, inability to rest and relax.
- **Social.** Risk taking, over- or under-eating, increased intake of alcohol or cigarettes, aggression, withdrawal, isolation.

#### *Psychosocial support*

- The term “psychosocial” refers to the dynamic relationship between the psychological and social dimensions of a person, where the dimensions influence each other. The psychological dimension includes emotional and thought processes, feelings and reactions. The social dimension includes relationships, family, community networks, social values and cultural practices.
- “Psychosocial support” refers to actions that meet the psychological and social needs of individuals, families and communities. Psychosocial support (PSS) requires training and supervision. Your supervisor can help you access the appropriate training before you begin to offer PSS to community members. They will also provide you with supervision and support while you provide PSS.
- We provide psychosocial support to help people who have been affected by a crisis. Volunteers should explain what psychosocial support is and if they are appropriately trained, they should offer to provide it to those who wish to receive it. Early and adequate psychosocial support can prevent distress and suffering from turning into more severe mental health problems.
- Psychosocial support during emergencies should ensure safety and promote calm, connectedness, personal and collective efficacy, and hope.



## What to do and how to do it

Psychosocial support activities include:

- Psycho-education
  - Explain how to identify signs of psychosocial distress
  - Provide advice on how to cope during outbreaks (e.g. maintaining a daily routine as much as possible; calling friends and family to keep in touch and show care for each other; fact-checking information about a disease against trustworthy sources)
  - Share tips about relaxation
- Health education can have a positive psychosocial impact:
  - Health education can help community members to better understand their health status, regain a sense of control and cope with their situation
  - While being ill, and even after medical clearance, it can be difficult for people suspected of infection to resume normal life. Educating communities about the nature of the disease, how it spreads – and does not spread – and how to protect against it is an important tool against fear and stigma
- Active listening: Ensure the affected population can raise their concerns, provide suggestions and feedback. This information is used to reduce fear, address rumours and misinformation and increase sense of agency and dignity of the affected population.
- Life skills and vocational skills activities/lessons.
- Creative activities, sports and physical activities.
- Restoring family links.
- Child friendly spaces.
- Supporting memorials and traditional burials.
- Support and self-help groups
  - These include efforts to help people in isolation or quarantine maintain contact with their relatives and friends.
  - Community volunteers that respond to crises are also exposed to loss, devastation, injury and death. It is therefore important to seek support from managers when needed, and to create a supportive environment by showing concern for staff and other volunteers.
- Psychological first aid



## 20. Isolating sick people

### Overview

- Some diseases are so dangerous or infectious that it is necessary to isolate people who are sick to prevent them from passing the infection to others. Isolation means separating those who are sick from those who are healthy.
- Isolating people is not imprisonment; it is not done forcibly and should be carried out with their consent. You need to explain to affected people and their families why isolation is necessary. People may also need to be isolated from their companion or livestock animal (and vice versa), if it brings the risk of transmission of disease. In extreme cases of progressive outbreaks, sick livestock animals may need to be culled (selectively slaughtered) to stop the spread of a disease to other animals and humans. Animal isolation, quarantine or culling should also not be done forcibly and should be carried out with their care-giver's consent.
- When a person is in isolation, the people who come into direct contact with them (for example, family members and healthcare providers) must be protected using the appropriate protective measures, like personal protective equipment (PPE). The number of people who are in direct contact with the sick person should be kept to a minimum. For example, only one family member should provide care, food, water, etc. to the sick person.

### What to do and how to do it

#### *Surveillance and detection*

1. Even if a disease requires isolation, you should continue monitoring, looking for cases of the disease (see Action Tool [Community-based surveillance](#)).
  - When cases are found, the need for isolation should be explained to those who are ill and their families.
  - Explaining isolation protects others, including members of the sick person's family. It may be easier for both the sick person and his or her family to understand why isolation is being proposed if they receive a complete explanation.
2. Provide support to the affected person, family members and healthcare providers to facilitate isolation.
  - This is not done forcibly, and the wishes of the family and sick person should be respected as much as possible.
  - Those who care for a person in isolation should be shown how to assist him or her and how to protect themselves. They should be provided with proper protective equipment.

#### *Management and care*

1. Provide psychosocial support to families (see Action Tool [Psychosocial support](#)). Isolation of a loved one can be very hard, for the family as well as the person in isolation.
2. Make sure that people in isolation have adequate food, water, and health care, as well as help to cope with any loss of livelihood.
3. Ask if those who need isolation have dependants (e.g. children, people with disabilities) that may need additional support. Inform your manager, who will liaise with the Red Cross Red Crescent protection

team for dependants to get the appropriate support.

### ***Social mobilization, messaging, and community engagement***

1. Talk to the family, community and elders/leaders to prevent stigma and social rejection.
  - Try to stop isolation from causing stigma or social rejection of sick people and their families.
  - Explain to the community why isolation is necessary.
  - Seek the help of elders and community leaders to fight stigma.



Talk to the community and elders to prevent stigma and social rejection.

## **Community messages**



24. Finding sick people

## 25. Mass vaccination campaigns

### Overview

- Outside of routine immunization schedules, vaccines may be provided in response to an outbreak through mass vaccination campaigns.
- A mass vaccination campaign occurs when the authorities give vaccinations to as many (appropriate) people as possible in a short period. Mass vaccinations take place in addition to routine vaccinations (see Action Tool [Routine vaccinations](#)). They may be organized because routine vaccinations cannot be given or in order to help control an epidemic. If necessary, this is also a good opportunity for volunteers to get vaccinated if they are eligible. Some examples of mass vaccination campaigns are: targeting children during a measles outbreak, mass vaccination campaigns during an Ebola outbreak or mass vaccination during the COVID-19 pandemic.
- Other types of vaccination campaigns target only specific at-risk groups. For example, “ring vaccination” during Ebola Virus Disease outbreaks targeting contacts and contacts of contacts.

### What to do and how to do it

#### ***Preparing to promote a mass vaccination campaign***

- Use the National Society's network to publicize outbreak response vaccination campaigns.
- Make sure you have all the relevant information and know where and when the vaccination campaign is taking place and who is supposed to be vaccinated.
- Find out the reasons for vaccination and the basic facts about the disease that vaccination will prevent.
- Under the advice of health professionals, inform the community of any expected side effects (especially common side effects) of the vaccine. This will help the community to know what to expect and to understand a normal response to the vaccine.

#### ***Promoting mass vaccination***

- Help health workers to ensure that all individuals at risk get vaccinated, including in hard-to-reach areas. Use simple and straightforward messages.
- Disseminate well in advance the location, vaccination site hours, number of vaccination days and age groups targeted.

#### ***Social mobilization, messaging and community engagement***

- The most important task is SOCIAL MOBILIZATION (see Action Tool [Social mobilization and behaviour change](#)).
  - Coordinate with the health authorities.
- Familiarize yourself with the habits and beliefs of members of your community and how they normally deal with vaccinations.
  - Meet community leaders and tell them about the campaign; get them to help reach the community.

- Talk to members of the community and explain how important vaccination is to protect their children.
- If some members of the community are afraid of vaccinations, assist community workers to calm and remove their fears.
  - Correct rumours and misinformation about vaccinations (see Action Tool, *Dealing with Rumours*).
- Use information, education and communication materials, such as the community message tools in this toolkit, because pictures always help people to understand messages better.
- If requested by health authorities and/or other actors responsible for administering the vaccines, support with: crowd control to manage the influx of people and the queue; assist in checking the target group age and characteristics; collaborate with community leaders to maintain order. Other tasks such as registering people, dispensing other items such as vitamins, or setting up vaccination spaces may be undertaken if requested by and in coordination with the health actor responsible for administering the vaccines.



Mass vaccination campaigns

## Community messages



16. Attending vaccination campaigns



15. Using vaccination cards



23. Encouraging healthy behaviours in a community

## 28. Physical distancing

### Overview

- Physical distancing is the practice of keeping people at a safe distance from each other during outbreaks of highly contagious diseases, to prevent diseases from spreading.
- This is different from isolation, because it is something that everyone in a community should practise, not just sick people. Physical distancing is an effective way to prevent contagious diseases from spreading.

### What to do and how to do it

#### *Encouraging physical distancing*

**Promote the use of physical distancing as a method of preventing the spread of disease.** It includes doing things like:

- Avoiding crowded places. For example, encourage people to:
  - Pray alone or in small groups, rather than in large groups
  - Do laundry near home rather than in communal areas
  - Go to markets or other crowded areas at a time of day when there are fewer people there
- Avoiding non-essential gatherings. For example, encourage people to:
  - Postpone weddings or similar events until safe to do so
  - Avoid festivals or community gatherings until it is safe to do so
    - Avoiding common greetings. For example, encourage people to avoid handshakes, hugging or kissing
    - Limiting contact with people at higher risk. For example, encourage people to avoid unnecessary close contact with young children or elderly people
- Keeping a safe distance from others when possible (the distance may change depending on the disease, ask for clarification from health authorities).


#### **Recognize there are times when social distancing may be difficult or impossible.**

- In situations when people are sharing overcrowded living spaces, such as displacement shelters, social distancing may not be possible. In cases like this it is important to: Promote the use (and, if possible, the distribution) of essential protective equipment, such as face masks.
- Social distancing may be more difficult for people working in certain sectors and locations, for example market vendors or domestic workers.
  - Target these groups to raise awareness about the modes of transmission of the disease and the protective measures that they can take to minimize their exposure
  - Advocate for safe working conditions and access to health services

**Be aware that social distancing can lead to isolation** and may create or worsen psychosocial problems for some people.



- See Action Tool [\*Psychosocial support\*](#) to determine how to help.



## Community messages



21. Physical distancing



## 34. Handwashing with soap

### Overview

Handwashing is one of the most important ways to prevent the spread of several epidemics, especially diarrhoeal diseases. Handwashing is easy and everyone (including children) can and should do it. To wash hands people must have access to water and soap.

Hands should be washed with soap:

- **BEFORE:**
  - Preparing food
  - Eating
  - Feeding a child
  - Breastfeeding
  - Caring for someone who is ill or treating a wound (yours or someone else's)
- **AFTER:**
  - Using the toilet
    - Men, boys, women and girls should wash their hands after using the toilet
    - Women and girls should engage in menstrual hygiene during their monthly menstrual cycles
      - Promote the use of clean, dry materials (disposable or reusable)
      - Promote changing menstrual materials and bathing as often as needed.
      - Discourage sharing reusable pads with anyone else
  - Cleaning a baby
  - Touching garbage or waste
  - Touching or feeding animals; handling raw meat
  - Blowing nose, coughing or sneezing
  - Treating wounds or caring for sick people
  - Coming into contact with a sick person in an epidemic (see Action Tool *Handwashing in a highly infectious epidemic*)

### What to do and how to do it

#### ***How to wash hands***

1. Wet your hands and rub them with soap.
2. Rub all parts of your hands together for 10 to 15 seconds.
3. Use lots of force (push your hands together hard) and remember to wash all surfaces, including the backs of the hands and between the fingers.


4. Rinse hands well so they are free of soap.
5. Dry hands with a paper towel. If there is no towel, wave them in the air until they are dry.



## Community messages



08. Washing hands with soap



09. When to wash hands

## 35. Handwashing in a highly infectious epidemic

### Overview

- Handwashing is a vital source of protection in epidemics of highly infectious diseases such as Ebola, Marburg fever, Lassa fever, plague, MERS and monkeypox. It is possible to become exposed to these diseases and to fall sick when trying to help people. Highly infectious germs are spread very easily through body fluids and certain diseases can even be transmitted via the bodies of people who are dead.
- Very vigorous and comprehensive handwashing is needed to protect yourself. (See Action Tools: [\*Volunteer protection and safety\*](#); and [\*Personal protective equipment \(PPE\) for highly infectious diseases\*](#))

### What to do and how to do it

#### ***When to wash hands in a highly infectious situation***


Hands should be washed with soap and/or disinfectant:

- BEFORE: preparing food, eating, feeding a child, breastfeeding, caring for someone who is ill, treating a wound (yours or someone else's)
- AFTER: using the toilet or cleaning a baby; touching garbage or waste; touching or feeding animals; handling raw meat; blowing nose, coughing or sneezing; treating wounds or caring for sick people; coming into contact with a sick person in an epidemic
- IN ADDITION: During an epidemic (of a highly infectious disease, or another disease such as diarrhoea or cholera, or a respiratory infection), it is a good idea to use this handwashing method every time you have touched something that can pass on an infection.

#### ***How to wash hands in highly infectious epidemics***

When working in highly infectious epidemics, it is critical to use all available forms of protection. One of these is handwashing. To wash hands correctly during a highly infectious epidemic, you must:

- Use soap or an alcohol-based disinfectant.
- Wash your hands as usual.
- Then wash thoroughly between your fingers.
- Scrub the tips of the fingers of the two hands together.
- Wash each of your thumbs with the other hand.
- Scrub the tips of the fingers of each hand on the palm of the other hand.



## Community messages



10. Steps for washing hands in epidemics

## 41. Handling and slaughtering animals

### Overview

- Animals can carry and spread germs that cause diseases. These include livestock animals such as cattle, domestic poultry birds, and pigs, companion animals such as dogs and cats, household vectors such as rats, and wild animals such as bats and wild birds.
- All animals can carry germs in their bodies when they are alive and in their meat and offals when they are slaughtered for food. Animal by-products, such as milk or eggs, can also carry germs.
- Animals that have died (including fetuses that have been aborted or animals that died during birth) can also carry germs.
- To avoid spreading diseases, it is important to wear personal protective equipment and follow recommended hygiene practices when dealing with live or dead animals and their by-products.

### Diseases transmitted by animals

Disease	Animals	Transmission
Rift Valley fever	Sheep, cattle, other animals. Signs and symptoms in infected animals: abortion; deaths in young.	Transmitted by mosquito bites; droplets in the air; the blood of sick animals; the bodies of infected dead animals.
Avian influenza	Wild and domestic poultry (birds). Signs and symptoms in infected animals: death or neurological signs; may have no signs of illness.	Transmitted through droplets in the air; feathers; and potentially the eggs and meat of infected birds.
Mpox	Mainly monkeys. Also rats, squirrels and prairie dogs. Signs and symptoms in infected animals: mainly none. Skin sores, breathing problems in prairie dogs.	Transmitted by touching infected animals or their body fluids; the bite or scratch of an infected animal; the meat of infected animals.
Plague	Mainly rats. Also rabbits, squirrels and prairie dogs. Signs and symptoms in infected animals: none in above. May cause disease in cats and occasionally dogs.	Transmitted by flea bites and domestic rats; droplets in the air; the bodies of dead infected animals.

Disease	Animals	Transmission
<b>Leptospirosis</b>	Cattle, pigs, rodents (mainly rats). Signs and symptoms in infected animals: abortion; liver and kidney disease. None in rats.	Transmitted by contact (through a cut, eyes, mouth etc.); or ingestion of the urine of an infected animal.
<b>MERS</b>	Camels. Signs and symptoms in infected animals: none.	Transmitted by close contact with an infected animal or person.
<b>Hantavirus pulmonary syndrome</b>	Rodents (mainly rats). Signs and symptoms in infected animals: none.	Transmitted by droplets of rodent urine or faeces in the air; the bodies of dead infected rodents; a bite or a scratch.
<b>Anthrax</b>	Sheep, cattle (cows) and other animals. Signs and symptoms in infected animals: sudden death in sheep and cattle; neck swelling with breathing difficulties in pigs, dogs and cats.	Transmitted by contact with, or eating, infected animal products; spores in the air.
<b>Rabies</b>	Primarily dogs, but may affect other mammals such as cats, livestock and wildlife; initial signs include fever, pain and unusual or unexplained tingling, pricking, or burning sensations at the wound site, later progressive and fatal neurological signs	Dog bites and scratches cause 99% of the human rabies cases

### *Handling poultry and livestock*

Poultry (birds)	Sheep and cows
<ul style="list-style-type: none"> <li>• Be careful when keeping poultry (birds) at home: keep the birds separated from the places where humans live and ensure that where they live is regularly kept clean and safe.</li> <li>• Keep domestic poultry birds away from the reach and contact of wild birds.</li> <li>• Keep the birds in a closed yard or enclosed free-range area where they have plenty of room to roam.</li> <li>• Do not let children play where the birds are kept.</li> <li>• Cook poultry (meat or eggs) thoroughly.</li> <li>• When preparing raw meat or eggs, always use clean utensils (knives, forks) and clean surfaces.</li> <li>• After cooking, use new utensils for eating. Never use the same utensils that you cook with. If no unused utensils are available, wash the utensils thoroughly before they are used for eating.</li> <li>• When slaughtering and handling poultry, be sure to protect yourself. Wear gloves and goggles if possible.</li> <li>• Never touch sick or dead birds without protection.</li> <li>• Observe birds for signs of sickness. If you detect illness, inform the animal health and welfare authorities.</li> <li>• At the slaughterhouse, abattoirs or when slaughtering at home, animals with visibly infected tissue and organ parts (e.g. tubercles, worms, eggs of parasites, etc.) should be immediately condemned and disposed of/destroyed. They should not be consumed or taken to the market for sale.</li> <li>• Wash hands thoroughly with soap and water after any contact with living or dead animals (grooming, feeding, husbandry, etc.), their by-products (eggs, feathers, raw wool, etc.).</li> <li>• Dispose of animal carcasses appropriately and swiftly (by burning or burying).</li> </ul>	<ul style="list-style-type: none"> <li>• Keep sheep or cows in an enclosed yard or free range where they have plenty of room to roam.</li> <li>• Do not let children play where animals are kept.</li> <li>• Cook meat thoroughly.</li> <li>• Boil milk before drinking or cooking it.</li> <li>• When preparing raw meat, always use clean utensils (knives, forks) and clean surfaces.</li> <li>• After cooking, use new utensils for eating. Never use the same utensils that you cook with. If no unused utensils are available, wash the utensils thoroughly before they are used for eating.</li> <li>• When handling or slaughtering livestock, be sure to protect yourself. Wear gloves and goggles if possible.</li> <li>• Never touch sick or dead sheep, goat or cows without protection.</li> <li>• Observe animals for signs of sickness. If you detect illness, inform the animal health and welfare authorities.</li> <li>• At the slaughterhouse, abattoirs or when slaughtering at home, animals with visibly infected tissue and organ parts (e.g. tubercles, worms, eggs of parasites, etc) should be immediately condemned and disposed of/destroyed. They should not be consumed or taken to the market for sale.</li> <li>• Wash hands thoroughly with soap and water after any contact with living or dead animals (grooming, feeding, husbandry, etc.), their by-products (eggs, feathers, raw wool, etc.).</li> <li>• Dispose of animal carcasses appropriately and swiftly (by burning or burying).</li> </ul>

### ***Handling bushmeat (monkeys, rodents, wild pigs and others)***

Bushmeat should be discouraged as a food source, especially in areas where epidemic-causing infections are present in bushmeat. However, sometimes people may continue to eat bushmeat and to use bushmeat by-products. In these cases, volunteers should promote the following messages:

- Do not let children play where animal carcasses are kept.
- Cook bushmeat thoroughly.




- When preparing raw meat, always use clean utensils (knives, forks) and clean surfaces.
- After cooking, use new utensils for eating. Never use the same utensils that you cook with.
- When slaughtering, butchering or skinning bushmeat, be sure to protect yourself. Wear gloves and goggles if possible.
- If you come upon an animal in the bush that appears sick or is already dead, do not handle it without protection (for example, gloves).
- Observe animals for signs of sickness. If you detect illness, inform the authorities.
- Wash hands thoroughly with soap and water after any contact with living or dead animals.
- Dispose of animal carcasses appropriately and swiftly (by burning or burying).

## Community messages



## 08. Washing hands with soap



## 09. When to wash hands



25. Handling and slaughtering animals

## 42. Promoting safe sex

### Overview

Some epidemic-causing diseases can be passed from one person to another during sex, usually in semen, vaginal fluids or blood. Some of these diseases can continue to be spread through sex, even after the person has recovered from the disease. "Sex" means any kind of oral, vaginal or anal sex, or sharing of sex toys. Some epidemic-causing diseases that are known to be passed through sex are:

- Zika virus
- Ebola virus disease
- Marburg fever
- Lassa fever
- Mpox virus

There are many other diseases, such as HIV/AIDS, which can be passed through sexual activities. Promoting safer sex is an important public health message that can save many lives. Having "safer sex" means protecting yourself and your partner from diseases that can be transmitted during sexual activity.

### ***When to promote safer sex?***

During outbreaks of Zika, Ebola, Marburg fever, mpox or Lassa fever, it is important to include messaging about safer sex. Messages should include:

- How the disease is transmitted through sex
- How to practise safer sex

While safer sex messaging is very important to prevent the spread of epidemic disease, it is only one way that community members should protect themselves from becoming ill during outbreaks of the above diseases. Especially in the cases of Ebola, Marburg, mpox and Lassa fevers, the diseases are also spread through other close contact with infected people, not just sex. Practising safer sex alone will not protect people from these diseases. Safer sex is only one way to prevent disease. It should not be the principal focus of your messages during an epidemic.

### ***Messages about safer sex during epidemics should promote:***

- Use of male or female condoms when having sex.
  - Condoms are a barrier and block any infectious semen or fluid that may transmit an infection.
- Other ways (instead of sexual intercourse) of sharing intimacy with sexual partners

### ***Key facts about Zika:***

- A pregnant woman can pass Zika to her unborn baby which can cause severe brain defects in the baby
  - Pregnant women who live in or travel to places where Zika is present, or whose sexual partners live in or travel to places where Zika is present, should be instructed to go to a health centre for a check-up and to discuss Zika risk with their healthcare provider.

- In these cases, it is best not to have sex during pregnancy, or to use condoms during the whole pregnancy.
- If women are planning a pregnancy and live in an area where Zika is present, it is important to discuss the risks, protect both partners from mosquito bites, and consider postponing pregnancy until after the outbreak has ended
- A man or woman who has Zika can pass Zika to his or her partner during sex for up to six months after acquiring the infection (whether they were sick and showed symptoms or not).

### ***Key facts about Ebola, Marburg and Lassa:***

- Men who have recovered from Ebola, Marburg fever or Lassa fever can pass the disease on to another person during oral, vaginal or anal sex
- Men who have recovered from Ebola or Marburg should use a condom for at least 12 months from when they got sick or until their semen tests negative twice for the virus.
- Men who have recovered from Lassa fever should use a condom for at least three months after they get better.

### ***Key facts about mpox***


- During the global outbreak that began in 2022, the virus mostly spread through sexual contact. Anyone who has close contact with someone who has symptoms and any person with multiple sex partners is also at risk. Therefore, people who have multiple or new sexual partners including those who are sexually active or gay, bisexual and other men who have sex with men may be at higher risk of being exposed if they have sex or other form of close contact with someone who is infectious.
- Therefore, it is important to engage communities including those of sex workers, gay, bisexual and other men who have sex with men to raise awareness on risks of mpox.

## **What to do and how to do it**

### ***Social mobilization, messaging, and behaviour change***

1. Make sure you understand the facts, and how diseases including Zika, Ebola, Marburg fever, mpox and Lassa fever can be transmitted through sex.
2. Remember that sex is only one way of transmitting these diseases, and often it is not the main way.
  - Talk to your volunteer coordination/team leader or local branch health office about which prevention methods should be emphasized during your community visits.
3. Remember that sex can be a very sensitive, taboo or embarrassing topic for people to talk about. You may have to change your approach or way of communicating to get your messages across.
4. Carry out social mobilization and behaviour change communication activities in an outbreak of Zika, Ebola, Marburg fever, mpox or Lassa fever. (See Important points above and Action Tools [\*Communicating with the community\*](#) and [\*Social mobilization and behaviour change\*](#).)
  - Make sure you know and can demonstrate how to use a male and female condom correctly.
5. Be respectful of the culture:
  - Consider speaking to men and women separately

- Consider speaking separately to adolescents. Remember that adolescents often experience stigma and difficulties in accessing sexual health information and services. Yet, most people initiate sexual activity during adolescence, so it is important to support them to ensure good sexual health choices and decisions.
- Consider adapting specific communication strategies and schedules to key populations as they are often marginalized and even hard-to-reach due to stigmatization or unfavorable laws in some countries.
- Do not push people to share views on this topic in front of others as some may feel uncomfortable
- Do not make assumptions based on stereotypes. For example, do not assume a person has certain attitudes about sex based on their age, gender, profession, or ethnicity
- Remember that providing sexual health information is important, even if it is a sensitive topic in many cultures



## Community messages



26. Practising safe sex

## 43. Social mobilization and behaviour change

### Overview


There are many reasons why people practise unhealthy behaviours. People are affected by access to services or facilities, social norms and influences where they work, live or play. Behaviour change is the study of how and why people change some habit or action in their life. As volunteers, we need to understand WHY the behaviour is happening and WHAT actions will lead to change to create healthy behaviours. Examples of healthy behaviours include handwashing, breastfeeding, immunizations, consistent condom use and use of bed nets.

In any culture and context, behaviour change involves three elements. Before people will change their behaviour:

1. They need to know what, why and how they should change. They need knowledge.
2. They need to have the right equipment, access and capacity. They need an enabling environment.
3. They need to be motivated to change.



The social-ecological model below shows how each person's behaviours are affected by many different levels of influence including the individual level, the interpersonal level, the community level, the organizational level and the broader policy level which includes laws and policies that allow or restrict a behaviour. In order to promote health, it is important to consider and plan behaviour change activities across multiple levels at the same time. This approach is more likely to result in successful behaviour change over time. As a volunteer, it is helpful to understand that behaviour change is difficult for many people because of these many levels and the complex interactions and expectations across the different levels. If you consider how each of the levels affects the behaviour of the person you want to help, you can try different interventions at each level that is specific to their needs.



Socio-ecological model

### What do to and how to do it

The general process for developing a behaviour change intervention includes staff and volunteers working through the general steps of:

1. Sensitizing the community to the behaviour change process using the theory of change model.
2. Assessing the problem behaviour – why it is practised, who practises it, when it is practised and what factors in the environment or society encourage the behaviour. Assess this information at the different levels of the social-ecological model for each community you serve.
3. Identifying an appropriate behaviour goal based on your assessment.
4. Reviewing the causes or barriers at each level that allow the behaviour to continue. Identify interventions

that align with each cause or barrier and that can be used at different levels.

5. Discussing the suggested interventions for each social-ecological model level with the community.
6. Identifying appropriate interventions for the context at each level. Interventions should be planned to address the stages of the theory of change by first giving knowledge and addressing environmental factors, motivating key people to gain approval and intentions, and ultimately inciting people to action that contributes to the overall goal.
7. Implementing the interventions at each level.
8. Monitoring to see if change is happening. Change takes time but it must be monitored to ensure that it is happening, even slowly. Additionally, as people go through the change process, their barriers and causes will change. The behaviour change interventions should adjust to these changes to ensure that change can continue.
9. Recognizing that when change is not happening as intended, further assessment and intervention tweaking is needed.
10. Continuing to implement, monitor, assess and adjust as the change process happens.

For more information, please consult the eCBHFA Manual for volunteers on [Behaviour Change](#), including:

1. Principles of behaviour change
2. The social ecological model
3. The stages of behaviour change
4. Activities for behaviour change

## Community messages



23. Encouraging healthy behaviours in a community